



**HAL**  
open science

## Balchit Obsidian (Upper Awash, Ethiopia)

Jean-Paul Raynal, Gérard Poupeau, Guy Kieffer, Marcello Piperno, J. Andy Milton, Sarah Delerue, François-Xavier Le Bourdonnec

► **To cite this version:**

Jean-Paul Raynal, Gérard Poupeau, Guy Kieffer, Marcello Piperno, J. Andy Milton, et al.. Balchit Obsidian (Upper Awash, Ethiopia). Colloque du GMPCA "Archéométrie 2005". INSTN-CEA, Saclay, 19-22 avril 2005., 2005. halshs-00004023

**HAL Id: halshs-00004023**

**<https://shs.hal.science/halshs-00004023>**

Submitted on 7 Jul 2005

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# BALCHIT OBSIDIAN (UPPER AWASH, ETHIOPIA)

Jean-Paul Raynal<sup>1</sup>, Gérard Poupeau<sup>2</sup>, Guy Kieffer<sup>3</sup>, Marcello Piperno<sup>4</sup>,  
J. Andy Milton<sup>5</sup>, Sarah Delerue<sup>2</sup>, François-Xavier Le Bourdonnec<sup>2</sup>




<sup>1</sup> Institut de Préhistoire et de Géologie du Quaternaire, UMR 5199 CNRS PACEA, Université Bordeaux 1, Avenue des Facultés, 33405 Talence, France et GDR 1122 CNRS. jpraynal@wanadoo.fr  
<sup>2</sup> Centre de Recherche en Physique Appliquée à l'Archéologie, IRAMAT, UMR 5060 CNRS-Université Bordeaux 3, Maison de l'Archéologie, Esplanade des Antilles, 33607 Pessac, France. Gerard.Poupeau@u-bordeaux3.fr, sarahdelerue@hotmail.com, françois.lebourdonnec@etu.u-bordeaux3.fr  
<sup>3</sup> Centre de Recherches Volcanologiques, UMR 6042 CNRS, Université Blaise Pascal, 29 boulevard Gergovia, 63037 Clermont-Ferrand Cedex 1, France, et GDR 1122 CNRS.  
<sup>4</sup> Dipartimento di Scienze Storiche, Archeologiche e Antropologiche dell'Antichità, Università di Roma "La Sapienza" et Directeur de la Mission archéologique italienne à Melka Kunture. m.piperno@tin.it  
<sup>5</sup> School of Ocean and Earth Science, Southampton Oceanography Centre, University of Southampton, Empress Dock, Southampton SO14 3ZH, United Kingdom. Jam2@soc.soton.ac.uk

The small volcanic massif of Balchit is one of the Pliocene silicic centers of the Wechecha Formation (Addis Ababa Rift Embayment). It is located some 25 km SE of Addis Ababa and 7 km NE of Melka Kunture, on the left interfluve of the Awash river (see map). One Balchit obsidian was recently dated at  $4.37 \pm 0.07$  Ma by K-Ar (Chernet *et al.*, 1998).



Accumulation of obsidian artefacts and debris.



## Obsidian and man in the Melka Kunture area

In the vicinity of Melka Kunture, obsidian is a major component of lithic series since the Oldowayan. The nearby Balchit volcanic massif constitutes a major obsidian source-area (Berthelet *et al.*, 2001). Since 1999, new investigations have been undertaken on the prehistory and volcano-sedimentary environments of Melka Kunture and a special attention was paid to obsidian artefacts and its primary and secondary sources. Analyses were performed on several obsidian samples from various locations, both *in situ* from a lava flow of Balchit and reworked debris or pebbles from different alluvial formations of the Awash river and its tributaries (Poupeau *et al.*, 2004).



Close-up view of obsidian artefacts.



Balchit volcanic complex: view from top towards Balchit village with Wechecha volcano in background.

## Obsidian occurrences

The only recognized source of obsidian is the Balchit flat flow-dome, which offers a few kilometres square outcrop with a wide variety of eruptive facies (vertical and convolute fluidal structures, finely banded perlitic lava, spherulitic facies, massive obsidian, etc). Remarkable amygdals up to metric lenses of pure and massive obsidian are scattered among the various lava facies. The obsidian colour is dominantly black but locally blue, green, red and beige colours have been observed.

As products of erosion, blocks, cobbles, pebbles and gravels are found in quaternary alluviums and in minor river beds and form secondary sources which were available for prehistoric groups (Kieffer *et al.*, 2004).




Convolute facies.



Vertical facies.



Unweathered massive block of obsidian.



Normalized abundance of trace elements in Melka Kunture obsidians. The norm selected is the composition of the primitive earth mantle as given by Sun and McDonough (1989).

## ICP-MS analysis and comments

The trace element contents of two Balchit samples from the main obsidian-bearing lava flow and of nine obsidian pebbles from alluvial deposits were determined by ICP-MS at LGCA (Grenoble) and at SOCFAC (Southampton), using the same experimental procedure (Barrat *et al.*, 2000).

Six obsidians pebbles were found to present the same (type A) elemental composition as the two Balchit obsidians and therefore might come from this mother-rock.

The three other obsidians present specific trace element systematics dubbed respectively as types B, C and D (diagram on left).

## Conclusions

Field observations show that the large size of Simbiro Acheulian obsidian artefacts found in the area studied excludes a raw material procurement from the local alluvial deposits. This preliminary work shows that at least four geochemically different sources could have been exploited by Ancient Man. One of these sources could be the Balchit main lava flow or one of its proximal concentrations of erosion products.

Thus in order to deepen our understanding of the obsidian procurement strategies of Ancient Man, the next step (in progress) will be to draw a comprehensive map of the potential obsidian sources in the vicinity of the Melka Kunture area and to establish their geochemical fingerprinting.

## References

BARRAT J. A., BLICHERT-TOFT J., GILLET Ph. and KELLER F., 2000, The differentiation of eucrites: the role of in-situ crystallization. *Meteoritics Planet. Sci.* **35**, 1087-1100.  
 BERTHELETA., CHAVAILLON J., BULGARELLI G. M. and PIPERNO M. (Eds.), 2001, *Melka Kunture, The guide*, Finiguerra Arti Grafiche, Lavello.  
 CHERNET T., HART W. K., ARONSON J. L. and WALTER R. C., 1998, New age constraints on the timing of volcanism and tectonism in the northern Main Ethiopian Rift-southern Afar transition zone (Ethiopia). *Journal of Volcanology and Geothermal Research*, **80**, 267-280.  
 POUPEAU G., KIEFFER G., RAYNAL J. P., MILTON A. and DELERUE S., 2004, Trace element geochemistry in Balchit obsidian (Upper Awash, Ethiopia) in Jean Chavaillon and Marcello Piperno (Eds.), "Studies on the Early Paleolithic site of Melka Kunture, Ethiopia", 103-110.  
 KIEFFER G., RAYNAL J. P. and BARDIN G., 2004, Volcanic markers in coarse alluvium at Melka Kunture (Upper Awash, Ethiopia), in: Jean Chavaillon and Marcello Piperno (Eds.), "Studies on the Early Paleolithic site of Melka Kunture, Ethiopia", 93-101.

## Acknowledgments

We thank the Oromia Regional State and the Ministry of Youth, Sports and Culture of Ethiopia (Research and Conservation of Cultural Heritage and Department of Anthropology and Archaeology).