



HAL
open science

Dynamic equivalence principle in linear rational expectations models

Stéphane Gauthier

► **To cite this version:**

Stéphane Gauthier. Dynamic equivalence principle in linear rational expectations models. *Macroeconomic Dynamics*, 2003, 7 (3), pp.63-88. 10.1017/S1365100502010301 . halshs-00069499

HAL Id: halshs-00069499

<https://shs.hal.science/halshs-00069499>

Submitted on 12 Sep 2012

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

On the Dynamic Equivalence Principle in Linear Rational Expectations Models *

Stéphane Gauthier †

CREST, laboratoire de macroéconomie
and ERMES, Université Paris 2

2 Juillet 2000

Abstract

Linear models with infinite horizon admit in general infinitely many rational expectations solutions. Some additional selection devices are consequently needed in order to narrow the set of relevant solutions. The viewpoint of this paper is that a solution will be more likely to arise if it is locally determinate, that is, locally isolated, locally immune to sunspots, and locally stable under learning. These three criteria are applied to solutions of linear univariate models along which the level of the state variable evolves through time. In such models the equilibrium behavior of the level of the state variable is described by a linear recursive equation characterized by the set of its coefficients. The main innovation of this paper is to define a new perfect foresight dynamics whose fixed points are these sets of coefficients, thus allowing us to study the property of determinacy of these sets, or, equivalently, of the associated solutions. It is shown that only one solution is locally determinate in this new dynamics. It is also locally immune to sunspots and locally stable under myopic learning. This solution corresponds to the saddle path in the saddle point case.

JEL Classification Number: E32

Keywords: rational expectations, determinacy, sunspots, learning.

* This work has been undertaken at DELTA (Joint Research Unit CNRS-EHESS-ENS) and GREQAM (Université Aix-Marseille 2). I wish to thank Russell Davidson, Gabriel Desgranges, Jean-Michel Grandmont, Roger Guesnerie and two anonymous referees of this journal for many helpful comments. Of course the usual disclaimers apply.

† Stéphane Gauthier, CREST-LMA, bâtiment de Malakoff 2 (timbre J360), 15 bd Gabriel Péri, 92245 Malakoff cedex, France ;Phone number: 33 1 41 17 37 38; Fax number: 33 1 41 17 76 66; e-mail address: gauthier@ensae.fr.

1 Introduction

Linear models with infinite horizon admit in general infinitely many perfect foresight solutions. This implies in particular that there are infinitely many ways the economy may react to unanticipated shocks. Some claim that this makes such models useless for economic predictions or policy evaluations. Some even claim that non-uniqueness undermines the very concept of perfect foresight itself since there is then no clear reason why economic agents should manage to choose one particular solution (Kehoe and Levine 1985). The purpose of this paper is to describe three alternative conditions that should be met for public opinion to focus so sharply. Following Guesnerie 1993, we shall argue that a solution should be locally determinate, locally immune to sunspots, and locally stable under learning in order to be a possible outcome of a decentralized process through which agents try to coordinate their behavior on perfect foresight. The first two criteria are based on local multiplicity properties, as they respectively rule out any solution arbitrarily close to which there exists some other perfect foresight solution, or some stationary stochastic sunspot solution along which the system fluctuates in response to random events unrelated to economic fundamentals. In contrast, the last criterion recommends eliminating any solution that economic agents would fail to discover through a simple adaptive learning process. These three criteria have clearly very different status. It will be shown, however, that in the field of models covered in this paper they select the same solution, the one commonly referred to as the fundamental solution in the literature (McCallum 1983).

In nonstochastic models these three criteria have been primarily applied to steady states, i.e., particular trajectories along which the level of the state variable remains constant over time (see, e.g., Azariadis 1981, Azariadis and Guesnerie 1982, Chiapori, Geoffard and Guesnerie 1992, Farmer and Woodford 1997, Grandmont 1998, Grandmont and Laroque 1991, Guesnerie and Woodford 1991, Marcet and Sargent 1989 or Woodford 1984).¹ As emphasized by Guesnerie 1993, if economic agents forecast one period ahead and if there is no predetermined variables in the model, then a steady state is locally determinate if and only if it is locally immune to sunspots and locally stable for some reasonable learning rules. This suggests that these three criteria may consequently be equivalent, at least when attention is focused on a certain class of solutions to rational expectations models. This paper shows that this equivalence, suitably reinterpreted, may be viewed as a part of a *dynamic equivalence principle*, a property which provides, when it holds, a rather compelling selection device on the set of rational expectations equilibria of general linear models where there are arbitrarily many leads in expectations or lags in memory.

¹ In linear models subject to extrinsic shocks, there is a substantial literature on the stability under least squares learning of various dynamic rational expectations solutions. See, e.g., Evans and Honkapohja 1999 for a survey on this topics.

The relevant class of solutions to which this principle applies makes the current state of the system linked with the same number of past states as the number of initial conditions to the economic system (that is in turn equal to the number of predetermined variables). An appealing characteristic of these *minimal state variable solutions*² is that they can be fully defined by the set of the coefficients of the linear difference equation which governs the intertemporal behavior of the system in equilibrium. The main innovation of this paper is to apply the three criteria described above to these sets of coefficients in order to choose among minimal state variable solutions. Namely we shall study whether these sets of coefficients can be locally determinate, in a new dynamics with perfect foresight induced by the usual dynamics on the level of the state variable, locally free of finite Markovian stationary sunspot equilibria, and locally stable under a specified myopic learning process which fits the iterative version of the expectational stability device, extensively used by, e.g., DeCanio 1979 or Evans 1985. Surprisingly, it turns out that these three devices select the same minimal state variable solution, the so-called fundamental one. In particular, this solution coincides with the saddle path trajectory in the saddle point configuration, in accordance with the recommendations of the main selection devices used in the literature, such as the stability based device by Blanchard and Kahn 1981, the minimal state variable criterion by McCallum 1983, or the minimal variance criterion by Taylor 1977. Unlike these devices, however, this principle does not depend on particular features of the model, e.g., some stability properties, and there are quite natural reasons why the economy should apply it.

The paper is organized in the following way. Section 2 examines the relevance of the dynamic equivalence principle to the simple class of models with only one lead in expectations and one lag in memory. Section 3 reviews some of the results obtained by Desgranges and Gauthier 2001 and Gauthier 2001 in the more general class of one-step forward looking models where the number of lags in memory is arbitrary. Section 4, which contains the new results of the paper, is a preliminary exploration of the case where agents forecast beyond the next period, but where there is only one lag memory. Finally Section 5 concludes.

² See McCallum 1983 for a closely related terminology. McCallum 1983 actually defines minimal state variable solutions using both a primary principle and a second principle which isolates a particular minimal state variable solution as bubble-free. Here, as in Evans and Honkapohja 1999, we define minimal state variable solutions using only the primary principle. It is worth noticing that the dynamic equivalence principle recommends in fact to choose the solution identified as bubble-free, or fundamental, by McCallum 1983.

2 A Simple Framework

Let us first focus attention on the class of linear models where the current state of the economic system is a real number, e.g., a price, that depends on both the previous state and the forecast of the next state (we assume that this forecast is the same for all agents). Such a class encompasses some overlapping generations models with production, and some infinite horizon models with cash-in-advance constraints. Unfortunately, even in a so simple framework, there are multiple minimal state variable (hereafter MSV) solutions, and our purpose is to study whether some of them can be locally determinate, immune to sunspots, and stable under some adaptive learning rules. In general, in much of the literature, these devices have been primarily applied to steady states. Here, however, MSV solutions involve deterministic changes in the level of the state variable over time. Each of these solutions actually expresses the current state of the economic system as a linear function of the previous state. Therefore each of them can be fully characterized by a single coefficient, giving the constant ratio of two consecutive realizations of the level of the state variable, i.e., the growth rate of the level of the state variable. In this section, we shall apply the dynamic equivalence principle to each of these growth rates, or, equivalently, to the corresponding MSV solutions.³

Let the current state of the system be represented by a real number x_t at time $t \geq 0$ that depends on the past state x_{t-1} and on the common forecast $E(x_{t+1} | I_t)$ conditional to the information set I_t held by economic agents at time t , through the temporary equilibrium relation

$$\gamma_1 E(x_{t+1} | I_t) + x_t + \delta_1 x_{t-1} = 0, \quad (1)$$

where γ_1 and δ_1 are real parameters. Perfect foresight solutions to (1) are sequences (x_t) associated with the initial condition x_{-1} and satisfying the difference equation

$$\gamma_1 x_{t+1} + x_t + \delta_1 x_{t-1} = 0, \quad (2)$$

which results from replacing the forecast $E(x_{t+1} | I_t)$ by the actual realization x_{t+1} in (1). We shall restrict our attention to the particular class of MSV solutions to (2), which relate by definition the current state of the system to as many lagged variables as the number of initial conditions to (2). Therefore, along any of these solutions, the current state x_t will only depend on the previous state x_{t-1} through a law of motion of the form

$$x_t = \beta x_{t-1}. \quad (3)$$

³ In fact local stability under least squares learning of MSV solutions has been extensively considered in the literature. See, e.g., Evans and Honkapohja 1999 for a survey. Whether these solutions can be locally determinate or locally free of sunspots, however, has never been studied so far.

Following Grandmont and Laroque 1991, the growth rate β can be obtained by interpreting (3) as the outcome of a process where agents *a priori* believe that the economic system will evolve according to (3) whatever x_{t-1} and t are, and where this belief is self-fulfilling. According to this interpretation, traders form their forecast by leading (3) forward and set $E(x_{t+1} | I_t)$ equal to βx_t in (1). Thus, the actual law of motion corresponding to this forecast, directly obtained from (1), yields

$$x_t = -(\delta_1/(1 + \gamma_1\beta))x_{t-1}, \quad (4)$$

provided that $\beta \neq -1/\gamma_1$. It is clear that the actual law (4) coincides with the *a priori* guess (3), for any x_{t-1} and any t , if and only if the actual growth rate in (4) is equal to the *a priori* guess on the growth rate in (3), or, equivalently,

$$-(\delta_1/(1 + \gamma_1\beta)) = \beta \Leftrightarrow \gamma_1\beta^2 + \beta + \delta_1 = 0. \quad (5)$$

Let λ_1 and λ_2 be the roots of (5). Assume that they are *real* and *distinct*, i.e., $1 - 4\gamma_1\delta_1 > 0$. Let also λ_1 be the root of least modulus, i.e., $|\lambda_1| < |\lambda_2|$. The process described in this section highlights the point that, given the previous state of the system, the occurrence of an MSV solution depends on whether traders will be able to choose the root λ_i ($i = 1, 2$) corresponding to it. As in Guesnerie 1993, we shall argue that economic agents may not succeed in coordinating their behavior on a root that fails to be locally determinate, immune to sunspots, or stable under some learning rule.

2.1 Determinacy of growth rates

According to the determinacy criterion, coordination on a root of (5) will be necessarily disturbed if there is at least another perfect foresight solution (x_t) to (2) that displays a sequence of growth rates ($\beta_t \equiv x_t/x_{t-1}$) arbitrarily close to this root. In order to assert whether some of these roots may be locally determinate, it must be noticed that the perfect foresight dynamics of the level of the state variable, described by (2), induces a new dynamics on the growth rate whose fixed points are the two roots λ_1 and λ_2 which solve (5). This new dynamics is obtained by assuming that the level of the state variable evolves according to (2) and is bound to satisfy in addition the relation

$$x_t = \beta_t x_{t-1} \quad (6)$$

whatever x_{t-1} and t are. It follows from (6) that (2) can be rewritten as

$$\gamma_1\beta_{t+1}\beta_t x_{t-1} + \beta_t x_{t-1} + \delta_1 x_{t-1} = 0,$$

since $x_{t+1} = \beta_{t+1}x_t$ in virtue of (6). Thus, if $x_{t-1} \neq 0$, i.e., if the economic system is not at the steady state level of the state variable, one can define the growth rate

perfect foresight dynamics induced by (2) as a nonlinear sequence of growth rates (β_t) such that

$$\gamma_1\beta_{t+1}\beta_t + \beta_t + \delta_1 = 0 \quad (7)$$

in each period $t \geq 0$. This dynamics is well defined if $\beta_t \neq 0$ and $\beta_t \neq -1/\gamma_1$ whatever t is. Its fixed points are the roots λ_1 and λ_2 . Hence (7) is locally well-defined if and only if $\lambda_i \neq 0$ ($i = 1, 2$), which holds true if both $\gamma_1 \neq 0$ and $\delta_1 \neq 0$ (since $\lambda_1 + \lambda_2 = -1/\gamma_1$ and $\lambda_1\lambda_2 = \delta_1/\gamma_1$). Observe now that the current growth rate β_t defined in (6), i.e., equal to the ratio x_t/x_{t-1} , is not predetermined in (7) at time t because x_t is not predetermined in (2) at this date. Therefore (7) has a classical one-step forward looking structure without predetermined variables, and the root λ_i is locally determinate in (7) if and only if it is locally unstable in the forward perfect foresight dynamics (7). The following result is due to Gauthier 2001.

Proposition 1 *Let $\gamma_1 \neq 0$ and $\delta_1 \neq 0$ in (1). Then, the root of lowest modulus (λ_1) is locally determinate in the growth rate perfect foresight dynamics (7), while the root of highest modulus (λ_2) is locally indeterminate in this dynamics.*

Proof. The dynamics (7) arbitrarily close to λ_i ($i = 1, 2$) is governed by the first order approximation of (7) at point $\beta_t = \beta_{t+1} = \lambda_i$ ($i = 1, 2$),

$$\gamma_1\lambda_i(\beta_{t+1} - \lambda_i) + (\gamma_1\lambda_i + 1)(\beta_t - \lambda_i) = 0. \quad (8)$$

The root λ_i is locally unstable in (8) if and only if $|(\gamma_1\lambda_i + 1)/(\gamma_1\lambda_i)| > 1$. Since $(\lambda_1 + \lambda_2) = -1/\gamma_1$, this inequality rewrites

$$|(\gamma_1\lambda_i + 1)/(\gamma_1\lambda_i)| = |(\lambda_i + 1/\gamma_1)/\lambda_i| = |\lambda_j/\lambda_i| > 1,$$

for $j \neq i$, $j = 1, 2$. Thus, the root λ_1 is locally unstable ($|\lambda_2/\lambda_1| > 1$) and the root λ_2 is locally stable ($|\lambda_1/\lambda_2| < 1$). *Q.E.D.*

Proposition 1 recommends to pick out the MSV solution corresponding to the root of least modulus, independently of the stability properties of the dynamics (2), i.e., independently of $|\lambda_1|$ and $|\lambda_2|$. However, if (2) is to be interpreted as a dynamics restricted to an arbitrarily small neighborhood of the steady state ($x_t = 0$), then the stability condition $|\lambda_1| < 1$ should hold for the corresponding MSV solution to be locally feasible. This condition covers both the saddle point configuration ($|\lambda_1| < 1 < |\lambda_2|$), where the locally determinate MSV solution coincides with the stable saddle path ($x_t = \lambda_1 x_{t-1}$), and the sink configuration ($|\lambda_2| < 1$) where there are infinitely many stable solutions to (2).

2.2 Sunspots on growth rates

Suppose now that traders form their expectations about the rate of change of the level of the state variable conditionally to a k -state Markovian sunspot process associated with a Markov matrix $\mathbf{\Pi}$. Note that traders are not directly concerned here with the level of the state variable itself, as is usually the case in the literature. Instead, if the current sunspot event s_t is s ($s = 1, \dots, k$), they *a priori* believe that $\beta_t = \beta_s$ and expect $\beta_{t+1} = \beta_{s'}$ to occur with probability $\pi_{ss'}$ ($s' = 1, \dots, k$). Their forecast, conditional to this sunspot event, writes then

$$E(x_{t+1} | \{x_t, s_t\}) = \left(\sum_{s'=1}^k \pi_{ss'} \beta_{s'} \right) x_t \equiv \bar{\beta}_s x_t, \quad (9)$$

where $\bar{\beta}_s$ is a one-period average growth rate. The actual dynamics, obtained by reintroducing the forecast (9) into the temporary equilibrium map (1), i.e.,

$$x_t = -(\delta_1 / (1 + \gamma_1 \bar{\beta}_s)) x_{t-1}, \quad (10)$$

is consistent with the *a priori* belief of traders whenever the actual growth rate in (10) is equal to the *a priori* guess about the growth rate in sunspot event s , or, equivalently,

$$-(\delta_1 / (1 + \gamma_1 \bar{\beta}_s)) = \beta_s \Leftrightarrow (1 + \gamma_1 \bar{\beta}_s) \beta_s + \delta_1 = 0. \quad (11)$$

Following Desgranges and Gauthier 2001, we shall define a stationary sunspot equilibrium on the growth rate as a vector $\boldsymbol{\beta} = (\beta_1, \dots, \beta_k)$ associated with $\mathbf{\Pi}$ such that (a) there are s and $s' \neq s$ such that $\beta_s \neq \beta_{s'}$ ($s, s' = 1, \dots, k$), and (b) the pair $(\boldsymbol{\beta}, \mathbf{\Pi})$ satisfies (11) for any s ($s = 1, \dots, k$). Condition (a) ensures that the growth rate truly changes according to sunspot events, and condition (b) makes traders' beliefs consistent with rational expectations. Observe that the vector $\boldsymbol{\beta} = (\lambda_i, \dots, \lambda_i)$ satisfies condition (b) whatever $\mathbf{\Pi}$ is, but violates (a). Thus there can exist stationary sunspot equilibria on the growth rate $(\boldsymbol{\beta}, \mathbf{\Pi})$ such that each component β_s of $\boldsymbol{\beta}$ stands arbitrarily close to λ_i only if condition (b) does not implicitly define $\boldsymbol{\beta}$ as a smooth function of $\mathbf{\Pi}$ at point $(\lambda_i, \dots, \lambda_i)$, i.e., only if the implicit function theorem does not apply at this point.

Proposition 2 *Let a root λ_i ($i = 1, 2$) be immune to sunspots if there do not exist stationary sunspot equilibria on the growth rate $(\boldsymbol{\beta}, \mathbf{\Pi})$ such that each component β_s ($s = 1, \dots, k$) of $\boldsymbol{\beta}$ stands arbitrarily close to λ_i . Then, if $\gamma_1 \neq 0$ and $\delta_1 \neq 0$ in (1), the root of lowest modulus (λ_1) is the only one to be immune to sunspots.*

Proof. This follows from standard results in the class of one-step forward looking models without predetermined variables. See, e.g., Chiappori, Geoffard and Guesnerie 1992, or Desgranges and Gauthier 2001 for precise statements. *Q.E.D.*

2.3 Learning growth rates

Let us now focus attention on the case where traders try to discover some MSV solution through an adaptive learning process. Their *a priori* belief is assumed to be still consistent with (3) but the parameter β in this equation is no longer necessarily equal to some root λ_1 or λ_2 of (5). It is instead estimated by β_t at time t . Conditionally to this belief, the forecast $E(x_{t+1} | I_t)$ is equal to $\beta_t x_t$ in (1), and, as (4) shows, the actual law of motion of the state variable corresponding to this forecast is

$$x_t = -(\delta_1/(1 + \gamma_1\beta_t))x_{t-1}, \quad (12)$$

as long as $\beta_t \neq -1/\gamma_1$. It is clear that the actual growth rate $-(\delta_1/(1 + \gamma_1\beta_t))$ in (12) will differ from the estimated growth rate β_t as long as β_t is not equal to either λ_1 or λ_2 . This spread between the actual and the estimated growth rates should urge agents to revise their estimate in the next period. We shall use here one of the simplest learning rules, the myopic learning rule, which recommends to set the new estimate of the growth rate equal to the actual growth rate at time t , i.e., $\beta_{t+1} = -(\delta_1/(1 + \gamma_1\beta_t))$, or,

$$\gamma_1\beta_t\beta_{t+1} + \beta_{t+1} + \delta_1 = 0. \quad (13)$$

Since the dynamics with learning (13) is merely the time mirror of the perfect foresight dynamics of growth rates (7), the local stability properties of its fixed points λ_1 and λ_2 are the reverse of those given in Proposition 1.⁴

Proposition 3 *Let $\gamma_1 \neq 0$ and $\delta_1 \neq 0$ in (1). Then the root (λ_1) is locally stable in the dynamics with myopic learning (13) while the root of highest modulus (λ_2) is locally unstable in this dynamics.*

Proof. Obvious from Proposition 1. *Q.E.D.*

The dynamic equivalence principle directly follows from Propositions 1, 2, and 3, provided that both expectations and past matter (i.e., $\gamma_1 \neq 0$ and $\delta_1 \neq 0$, respectively). Both conditions are actually necessary for there exist multiple MSV solutions. In particular, in the case where $\delta_1 = 0$ in (1), the steady state ($x_t = 0$) is the unique MSV solution to the model, under the additional regularity condition that $\gamma_1 \neq -1$. As emphasized by Guesnerie 1993, it is then locally determinate in the usual dynamics with perfect foresight on the level of the state variable, i.e., (2) with $\delta_1 = 0$, if and only if it is locally immune to Markovian stationary sunspot equilibria on the level of the state variable, and locally stable under a myopic learning rule.

⁴ The algorithm (13) obviously belongs to a special class of learning rules. It is in fact identical to the iterative version of the expectational stability device used by DeCanio 1979 and Evans 1985. As shown by Evans and Guesnerie 1993, stability of a rational expectations solution in (13) is necessary, but not sufficient, for this solution to be strongly rational. Note also that if a solution is locally stable in the learning dynamics (13), then it is also locally expectationally stable in the sense of Evans 1989, and accordingly locally stable for the least squares learning rule, as first shown by Marcet and Sargent 1989.

3 One-Step Forward Looking Models

We shall now study whether the dynamic equivalence principle still holds true if $L \geq 1$ predetermined variables are introduced into the temporary equilibrium map (1). The evolution of the level of the state variable along any MSV solution is then described by an L -dimensional linear difference equation. Hence, given the L initial conditions to the economic system, a solution of this type will be no longer characterized by a single coefficient, a constant growth rate, but instead by the set of the L coefficients of this linear difference equation, to be called a *stationary extended growth rate*. This section describes how to apply the three tests used in Section 2 to such sets of coefficients, or, equivalently, to the corresponding MSV solutions. As shown by Gauthier 2001, only one MSV solution is locally determinate in the suitable dynamics with perfect foresight, whose fixed points are now these stationary extended growth rates. Moreover, as shown by Desgranges and Gauthier 2001, this solution is the only one to be locally immune to sunspots. Finally, as the local stability properties of this solution in the dynamics with myopic learning obtain again as a simple corollary of its property of local determinacy, the form taken here by the dynamic equivalence principle is still satisfied in the more general framework under consideration, and, furthermore, still allows us to pick out a unique MSV solution, the fundamental one.

Let the temporary equilibrium relation (1) be transformed as

$$\gamma_1 E(x_{t+1} | I_t) + x_t + \sum_{l=1}^L \delta_l x_{t-l} = 0. \quad (14)$$

Perfect foresight solutions are now sequences (x_t) associated with the initial condition (x_{-1}, \dots, x_{-L}) and satisfying the $(L + 1)$ -dimensional difference equation

$$\gamma_1 x_{t+1} + x_t + \sum_{l=1}^L \delta_l x_{t-l} = 0, \quad (15)$$

obtained by replacing the forecast $E(x_{t+1} | I_t)$ by the actual realization x_{t+1} in (14). This dynamics is governed by the $(L + 1)$ roots $(\lambda_1, \dots, \lambda_{L+1})$ of the characteristic polynomial associated with (15). Let these roots be real and distinct, with $|\lambda_1| < \dots < |\lambda_{L+1}|$.⁵ Unlike Section 2, however, we are not primarily concerned with these roots since the presence of $L \geq 1$ predetermined variables in (14) makes now the current state x_t linearly related to the $L \geq 1$ past realizations $(x_{t-1}, \dots, x_{t-L})$ along any MSV solution. These solutions actually satisfy not only (15) but also, in addition,

⁵ Theorem 6 in Gantmacher 1966 (Ch.15, §9) gives the number of real and distinct roots of a polynomial as a function of its real coefficients. A simple necessary condition for all these roots to be real is that $\gamma_1 \delta_1 \leq 1$, $\delta_2 \leq \delta_1^2$, $\delta_1 \delta_3 \leq \delta_2^2$, \dots , $\delta_{L-2} \delta_L \leq \delta_{L-1}^2$ (see Du Gua-Huat-Euler theorem in Mignotte 1989).

the difference equation

$$x_t = \sum_{l=1}^L \beta_l^* x_{t-l}, \quad (16)$$

where the vector of coefficients $\beta^* \equiv (\beta_1^*, \dots, \beta_L^*)$ is called a *stationary extended growth rate*. As in Section 2, one can characterize such a vector by interpreting (16) as a situation where (16) is a self-fulfilling belief about the evolution of the economic system. Assume accordingly that traders believe that the law of motion of the level of the state variable is given by (16) for some set of coefficients $(\beta_1, \dots, \beta_L)$. In this case, they form their forecast by iterating (16) once,

$$E(x_{t+1} | \{x_t, \dots, x_{t-l+1}\}) = \sum_{l=1}^L \beta_l x_{t-l+1}, \quad (17)$$

so that the actual evolution of the level of the state variable, which results from reintroducing (17) into (14), determines the actual current state x_t as a function of past history, namely

$$x_t = - \sum_{l=1}^L [(\gamma\beta_{l+1} + \delta_l)/(1 + \gamma\beta_1)] x_{t-l}, \quad (18)$$

with the convention that $\beta_{L+1} \equiv 0$. The belief (16) is then self-fulfilling if and only if (16) and (18) coincide whatever t and x_{t-l} are, or, equivalently, $\beta_l = \beta_l^*$ for $l = 1, \dots, L$, with

$$\beta_l^* = - \sum_{l=1}^L [(\gamma\beta_{l+1}^* + \delta_l)/(1 + \gamma\beta_1^*)], \quad (19)$$

whatever l is ($l = 1, \dots, L$), and $\beta_{L+1}^* \equiv 0$. How the solutions $(\beta_1^*, \dots, \beta_L^*)$ to (19) depend on economic fundamentals, summarized here by the parameters γ_1 and δ_l in (14), is fully described in Gauthier 2001. For our purpose, it is sufficient to notice that (19) admits in fact $(L+1)$ different solutions, i.e., there are $(L+1)$ stationary extended growth rates to the model (16). Indeed, since (15) and (16) must be consistent (otherwise the sequence of realizations of the level of the state variable induced by (16) would not form a perfect foresight equilibrium), it must be the case that (16) restricts the level of the state variable to evolve in one L -dimensional eigensubspace of (15), as x_t actually relies on L lagged variables in (16). Let W_i be the L -dimensional eigensubspace of (15) spanned by the L eigenvectors associated with the L roots in the set \mathcal{L}_i of all the roots but λ_i ($i = 1, \dots, L+1$). Since the total number of different L -dimensional eigensubspaces of (15) is equal to $(L+1)$, there are $(L+1)$ different MSV solutions (each of these solutions governs the behavior of the level of the state variable in a given the L -dimensional eigensubspaces of (15)). Now it follows from (16) that each MSV solution is in turn uniquely defined by a given set of the coefficients $(\beta_1^*, \dots, \beta_L^*)$ solution to (19) and by the L initial conditions to the economic

system. As a result, there are also $(L + 1)$ different stationary extended growth rates to the model (16). In the sequel, these vectors of coefficients will be denoted $\beta^*(\mathcal{L}_i)$, where $\beta^*(\mathcal{L}_i)$ actually corresponds through (16) to the MSV solution which governs the behavior of the level of the state variable in W_i .

3.1 Determinacy of extended growth rates

Our aim is now to examine whether some of these stationary extended growth rates pass the tests used in Section 3. Here we start applying the determinacy criterion by noticing that the usual dynamics on the level of the state variable, that is, (15), triggers a new dynamics on the vectors $\beta(t) = (\beta_1(t), \dots, \beta_L(t))$ whose fixed points are the $(L + 1)$ stationary extended growth rates. In this new dynamics, a stationary extended growth rate $\beta^*(\mathcal{L}_i)$ is locally determinate whenever there is no other vector $\beta(t)$ remaining arbitrarily close to it in each period. Otherwise it is locally indeterminate. As shown in Gauthier 2001, the stationary extended growth rate $\beta^*(\mathcal{L}_{L+1})$ corresponding to the set \mathcal{L}_{L+1} of the L roots of lowest modulus is in fact the only one to be locally determinate.

The *extended growth rate perfect foresight dynamics* comes by assuming that the level of the state variable is bound to satisfy (15) and, further, to evolve according to

$$x_t = \sum_{l=1}^L \beta_l(t) x_{t-l}, \quad (20)$$

whatever $t \geq 0$ and x_{t-l} ($l = 1, \dots, L$) are. Perfect foresight requires that the belief (20) be self-fulfilling, i.e., consistent with the actual law of motion of the system. This actual law is derived by replacing $E(x_{t+1} | I_t)$ in (15) by

$$x_{t+1} = \sum_{l=1}^L \beta_l(t+1) x_{t+1-l},$$

which is obtained by iterating once (20). It is then straightforward to verify that the actual law corresponding to (20) is

$$x_t = - \sum_{l=1}^L \left[(\gamma_1 \beta_{l+1}(t+1) + \delta_l) / (1 + \gamma_1 \beta_1(t+1)) \right] x_{t-l}, \quad (21)$$

with the convention that $\beta_{L+1}(t+1) \equiv 0$. We are now in a position to define the extended growth rate perfect foresight dynamics. It is a nonlinear sequence of extended growth rates $(\beta(t))$ such that (20) coincides with (21) whatever t and x_{t-l} are, or, equivalently,

$$\beta_l(t) = -(\gamma_1 \beta_{l+1}(t+1) + \delta_l) / (1 + \gamma_1 \beta_1(t+1)), \quad (22)$$

for $l = 1, \dots, L$. Its fixed points $\beta(t) = \beta(t+1)$ are the $(L + 1)$ stationary extended growth rates (which solve (19)). These dynamics are well defined arbitrarily close to

any stationary extended growth rate if and only if all the roots λ_i ($i = 1, \dots, L + 1$) differ from 0, which is true if and only if $\delta_L \neq 0$ (see Gauthier 2001). Since (22) has a one-step forward looking structure without predetermined variables, a stationary extended growth rate will be locally determinate in (22) if and only if all the L eigenvalues of the Jacobian matrix that governs the forward dynamics (22) in its immediate vicinity, have modulus greater than 1.⁶

Proposition 4 *The stationary extended growth rate $\beta^*(\mathcal{L}_{L+1})$ which governs the state variable perfect foresight dynamics (15) restricted to the L -dimensional subspace corresponding to the L roots of lowest modulus ($\lambda_1, \dots, \lambda_L$) is the only one to be locally determinate in the extended growth rate perfect foresight dynamics (22).*

Proof. See Gauthier 2001. *Q.E.D.*

Although Proposition 4 is independent of the stability properties of the dynamics with perfect foresight on the level of the state variable, the MSV solution corresponding to $\beta^*(\mathcal{L}_{L+1})$ will be locally feasible if and only if the condition $|\lambda_L| < 1$ is satisfied, which encompasses both the saddle point configuration ($|\lambda_L| < 1 < |\lambda_{L+1}|$) for (15), where $\beta^*(\mathcal{L}_{L+1})$ supports the saddle path trajectory, and the sink configuration ($|\lambda_{L+1}| < 1$) for this dynamics, where there are infinitely many stable solutions.

3.2 Sunspot equilibria on extended growth rates

We now come to study whether $\beta^*(\mathcal{L}_{L+1})$ is also the only stationary extended growth rate to be locally immune to sunspots. Suppose consequently that agents observe a k -state Markovian process associated with a Markov matrix $\mathbf{\Pi}$, and believe that, in the sunspot event $s_t = s$ ($s = 1, \dots, k$), the current state should be linked to the L previous states according to the law

$$x_t = \sum_{l=1}^L \beta_l^s x_{t-l}, \quad (23)$$

where the current set of coefficients $\beta^s \equiv (\beta_1^s, \dots, \beta_L^s)$ is allowed to depend on the current sunspot event. The belief (23) is intended to hold whatever t and the past history (x_{t-1}, \dots, x_{t-L}) are. Therefore agents also deduce from the occurrence of the event $s_t = s$ that the next extended growth rate should be equal to $\beta^{s'}$ with probability $\pi_{ss'}$ ($s' = 1, \dots, k$), and their forecast can consequently be written as

$$E(x_{t+1} \mid \{s_t = s, x_t, \dots, x_{t-L+1}\}) = \sum_{l=1}^L \left(\sum_{s'=1}^k \pi_{ss'} \beta_l^{s'} \right) x_{t+1-l}. \quad (24)$$

⁶ Equivalently, a stationary extended growth rate is locally indeterminate in (22) if at least one eigenvalue of the Jacobian matrix that governs the *backward* perfect foresight dynamics corresponding to (22) arbitrarily close to this stationary extended growth rate falls outside the unit disk. See, e.g., Chiappori, Geffard and Guesnerie 1992, p. 1098.

In a rational expectations equilibrium, the *a priori* belief (23) must be self-fulfilling whatever s_t is. If $s_t = s$, the actual dynamics corresponding to (23) is obtained by reintroducing the forecast (24) into (14), which leads to

$$x_t = - \sum_{l=1}^L \left[(\gamma (\sum_{s'=1}^k \pi_{ss'} \beta_{l+1}^{s'}) + \delta_l) / (\gamma (\sum_{s'=1}^k \pi_{ss'} \beta_1^{s'}) + 1) \right] x_{t-l}, \quad (25)$$

with the convention that $\beta_{L+1}^{s'} \equiv 0$ whatever s' is. Thus, in view of (23) and (25), one can define a stationary sunspot equilibrium on extended growth rates as a kL -dimensional vector $\boldsymbol{\beta} \equiv (\boldsymbol{\beta}^1, \dots, \boldsymbol{\beta}^k)$ associated with $\boldsymbol{\Pi}$ such that (a) there exist l ($l = 1, \dots, L$) and $s \neq s'$ ($s, s' = 1, \dots, k$) such that $\beta_l^s \neq \beta_l^{s'}$, and (b) β_l^s ($s = 1, \dots, k$) satisfies

$$\beta_l^s = -(\gamma \sum_{s'=1}^k \pi_{ss'} \beta_{l+1}^{s'} + \delta_l) / (\gamma \sum_{s'=1}^k \pi_{ss'} \beta_1^{s'} + 1)$$

for $l = 1, \dots, L$. Condition (b) makes the *a priori* belief (23) self-fulfilling since it ensures that (23) and (25) coincide whatever s_t is. For the extended growth rate to fluctuate according to sunspots, condition (a) must additionally hold. Otherwise, i.e., if (b) holds true but (a) fails, then $\boldsymbol{\beta}^s = \boldsymbol{\beta}^{s'} = \boldsymbol{\beta}^*(\mathcal{L}_i)$ for any $s, s' = 1, \dots, k$, and i ($i = 1, \dots, L + 1$).

The next result, which bears on Theorem 3 in Chiappori, Geoffard and Guesnerie 1992, studies whether there exist stationary sunspot equilibria on extended growth rates arbitrarily close to a stationary extended growth rate $\boldsymbol{\beta}^*(\mathcal{L}_i)$, i.e., such that the l th component β_l^s ($l = 1, \dots, L$) of $\boldsymbol{\beta}^s$ is arbitrarily close to the l th component $\beta_l^*(\mathcal{L}_i)$ of $\boldsymbol{\beta}^*(\mathcal{L}_i)$ whatever s and l are. A stationary extended growth rate $\boldsymbol{\beta}^*(\mathcal{L}_i)$ is said to be locally immune to sunspots when this is not the case.

Proposition 5 *The stationary extended growth rate $\boldsymbol{\beta}^*(\mathcal{L}_{L+1})$ which governs the state variable perfect foresight dynamics (15) restricted to the L -dimensional subspace corresponding to the L roots of lowest modulus $\lambda_1, \dots, \lambda_L$ is the only one to be locally immune to sunspots.*

Proof. See Chiappori, Geoffard and Guesnerie 1992 for a general study and Desgranges and Gauthier 2001 for an application to stationary sunspot equilibria on extended growth rates. *Q.E.D.*

3.3 Learning extended growth rates

It follows from Propositions 4 and 5 that a stationary extended growth rate is locally determinate if and only if it is locally immune to sunspots. We shall show in this section that the dynamic equivalence principle is satisfied since a stationary extended

growth rate is locally stable under myopic learning if and only if it is locally determinate. Assume, therefore, that the *a priori* belief on the law of motion still fits (16) whatever $t \geq 0$ and x_{t-l} ($l = 1, \dots, L$) are, but that agents are no longer aware of the entire set of stationary extended growth rates, and try to learn them. Let $\beta_l(t)$ be the estimate of the l th component of some stationary extended growth rate at date t . Given this vector of estimates, the traders' forecast $E(x_{t+1} | I_t)$ writes

$$E(x_{t+1} | \{x_t, \dots, x_{t-L}\}) = \sum_{l=1}^L \beta_l(t) x_{t-l+1}. \quad (26)$$

The actual law of motion of the level of the state variable, namely

$$x_t = - \sum_{l=1}^L (\gamma \beta_{l+1}(t) + \delta_l) / (1 + \gamma \beta_1(t)) x_{t-l}, \quad (27)$$

is obtained by reintroducing the forecast (26) into (14). In the myopic learning dynamics, traders compare their initial estimate $\beta_l(t)$ to the actual l th coefficient ($l = 1, \dots, L$) given in (27), and revise this estimate according to the rule

$$\beta_l(t+1) = - \sum_{l=1}^L (\gamma \beta_{l+1}(t) + \delta_l) / (1 + \gamma \beta_1(t)), \quad (28)$$

for any l , and with the convention that $\beta_{L+1}(t) \equiv 0$. The algorithm (28) can be thought of as a learning rule updated in real time in which agents keep fixed their forecast rule until the implied actual law of motion coefficients could be learned from the data, i.e., after L periods typically.⁷ As discussed in Evans 1985, the learning rule (28) is then identical to the iterative version of the expectational stability; it belongs nevertheless to a rather special class of learning rules. As (19) shows, the fixed points of this dynamics are the $(L+1)$ stationary extended growth rates. A stationary extended growth rate is locally stable under learning if and only all the eigenvalues of the Jacobian matrix that governs (28) in its immediate vicinity, have modulus less than 1. But (28) is simply the time mirror of the extended growth rate perfect foresight dynamics (22). Thus, the property of local determinacy of a stationary extended growth rate in (22) is equivalent to its property of local stability in (28), which establishes that the dynamic equivalence principle holds for the class of models (14).

Proposition 6 *The stationary extended growth rate $\beta^*(\mathcal{L}_{L+1})$ which governs the state variable perfect foresight dynamics (15) restricted to the L -dimensional subspace corresponding to the L roots of lowest modulus $\lambda_1, \dots, \lambda_L$ is the only one to be locally stable in the dynamics with myopic learning (28).*

Proof. Obvious from Proposition 4. *Q.E.D.*

⁷ The updating (28) is feasible every L periods provided that the $L \times L$ matrix generated is non-singular (see Evans 1985).

4 One-Lag in Memory Models

So far only models where agents forecast one-period ahead have been considered. In this section, we shall explore the remaining polar configuration where the number of leads in expectations is arbitrary, as our analysis will be restricted to the simple case where only one predetermined variable matters. As shown in Section 2, models of this class admit multiple MSV solutions. Each one makes the current state of the economic system linearly related to the previous state only, and, therefore, each one can be characterized by a constant growth rate of the level of the state variable. The question is whether the dynamic equivalence principle, when applied to such growth rates, is saved in this new framework. It will be shown that only one MSV solution is still both locally determinate and locally immune to sunspots, but this solution is no longer necessarily locally stable under myopic learning.

Let us consider the temporary equilibrium relation

$$\sum_{h=1}^H \gamma_h E(x_{t+h} | I_t) + x_t + \delta_1 x_{t-1} = 0, \quad (29)$$

where $E(x_{t+h} | I_t)$ represents the forecast formed at date t about the level of the state variable in period $t+h$ ($h = 1, \dots, H$), and where x_{t-1} is predetermined at date t . A perfect foresight solution is a sequence (x_t) associated with the initial condition x_{-1} and satisfying the $(H+1)$ -dimensional recursive equation

$$\sum_{h=1}^H \gamma_h x_{t+h} + x_t + \delta_1 x_{t-1} = 0, \quad (30)$$

obtained by assuming that $E(x_{t+h} | I_t)$ is equal to x_{t+h} in (29) whatever $h \geq 1$ and $t \geq 0$ are. Let λ_i ($i = 1, \dots, H+1$) be the $(H+1)$ roots of the characteristic equation associated with (30). As in the previous sections, we shall assume that these roots are real, distinct, and labelled in the order of increasing modulus, i.e., $|\lambda_i| < |\lambda_j|$ whenever $i < j$ ($i, j = 1, \dots, H+1$). Since there is only one predetermined variable in (29), the law of motion of the level of the state variable along any MSV solution to (30) is governed by a first order difference equation that still fits (3) and, now, the perfect foresight dynamics (30). The constant growth rate β in (3) ensures perfect foresight when (3) is a self-fulfilling belief, i.e., when it coincides with actual observations generated by (29). In view of (3), the forecast of traders is given by

$$E(x_{t+h} | I_t) = \beta E(x_{t+h-1} | I_t) = \dots = \beta^h x_t, \quad \text{for } h = 1, \dots, H, \quad (31)$$

so that the actual dynamics, resulting from reinserting (31) into (29), writes

$$x_t = -(\delta_1 / (1 + \sum_{h=1}^H \gamma_h \beta^h)) x_{t-1}. \quad (32)$$

Thus, the belief (3) on the law of motion coincides with the actual law (32), whatever x_{t-1} and t are, if and only if

$$\beta = -(\delta_1 / (1 + \sum_{h=1}^H \gamma_h \beta^h)) \Leftrightarrow \sum_{h=1}^H \gamma_h \beta^{h+1} + \beta + \delta_1 = 0, \quad (33)$$

or, equivalently, β is a root λ_i ($i = 1, \dots, H + 1$) of the characteristic polynomial associated with (30).

4.1 Determinacy of Growth Rates

The sequence of growth rates induced by the perfect foresight dynamics on the level of the state variable is obtained whenever the relation $x_t = \beta_t x_{t-1}$ holds in (30) whatever t and x_{t-1} are, so that $x_{t+h} = \beta_{t+h} x_{t+h-1} = (\beta_{t+h} \cdots \beta_t) x_{t-1}$ for $h = 0, \dots, H$. Under these requirements, the dynamics (30) rewrites

$$\sum_{h=1}^H \gamma_h \prod_{j=0}^h \beta_{t+j} x_{t-1} + \beta_t x_{t-1} + \delta_1 x_{t-1} = 0, \quad (34)$$

or, provided that $x_{t-1} \neq 0$,

$$\sum_{h=0}^H \gamma_h \prod_{j=0}^h \beta_{t+j} + \delta_1 = 0, \quad (35)$$

with the convention that $\gamma_0 \equiv 1$. The dynamics (35) is well-defined if $\beta_t \neq 0$ whatever t is. Its fixed points are the $(H + 1)$ roots λ_i ($i = 1, \dots, H + 1$) of (33). Hence it is well defined arbitrarily close to λ_i if and only if $\lambda_i \neq 0$. This is the case whatever i is if and only if $\delta_1 \neq 0$. The dynamics (35) arbitrarily close to λ_i is governed by a linear recursive equation of order H , obtained by linearizing (35) at point $\beta(t + s) = \lambda_i$ whatever s is ($s = 0, \dots, H$), namely

$$\sum_{h=0}^H \sum_{j=h}^H \gamma_j \lambda_i^j (\beta_{t+h} - \lambda_i) = 0. \quad (36)$$

Since (35) determines the current growth rate β_t as a function of future growth rates β_{t+h} ($h = 1, \dots, H$), a fixed point λ_i of (35) is locally determinate if and only if all the H roots of the characteristic polynomial associated to (36) have modulus greater than 1.

Proposition 7 *Assume that $\gamma_H \neq 0$ for $H \geq 1$ and $\delta_1 \neq 0$. Then the root of lowest modulus (λ_1) is the only fixed point of the growth rate perfect foresight dynamics (35) to be locally determinate in this dynamics.*

Proof. See in appendix. *Q.E.D.*

4.2 Sunspots on Growth Rates

Assume now that all the agents believe that the growth rate of the level of the state variable is perfectly correlated to a k -state Markovian sunspot process whose probability law is described by a $k \times k$ Markov matrix $\mathbf{\Pi}$. Hence agents expect the current growth rate β_t ($t \geq 0$) to be equal to β_s ($s = 1, \dots, k$) whenever they observe at date t some sunspot signal s . In a stationary sunspot equilibrium on the growth rate, this belief must be consistent with the actual law of motion of the system. This section is devoted to study whether stationary sunspot equilibria of this type exist when all the growth rates β_s are bound to lie arbitrarily close to a given root λ_i ($i = 1, \dots, H + 1$). The root of lowest modulus (λ_1) will be shown to be the only one to be immune to sunspot fluctuations, thus making local determinacy equivalent to the lack of local stationary sunspot equilibria on the growth rate.

In order to set out the actual temporary equilibrium dynamics that makes sunspot beliefs self-fulfilling, one must specify how traders forecast future at any date, given their *a priori* belief on the law of motion of the economic system. When the sunspot signal s_t at date t is s , traders expect (i) the period t growth rate to be β_s and (ii) the growth rates in the subsequent periods to be distributed conditionally to the event $s_t = s$, i.e., the probability that $\beta_{t+h} = \beta_{s'}$ ($s' = 1, \dots, k$) is given by the ss' th element of $\mathbf{\Pi}^h$. Since (i) holds true at any date, the expectation $E(x_{t+h} | \{s_t = s\})$ is equal to $E((\beta_{t+h} \cdots \beta_{t+1})x_t | \{s_t = s\})$. What is important, therefore, is the probability law of the product $(\beta_{t+h} \cdots \beta_{t+1})$. The following lemma describes this probability law.

Lemma 8 *Let \mathbf{B} be the $k \times k$ diagonal matrix whose ss th element is β_s ($s = 1, \dots, k$), i.e., the growth rate that traders expect to occur in sunspot state s . Let $x_t(s)$ be the temporary equilibrium state in sunspot state s at date t ($t \geq 0$). Let \mathbf{X}_t be the $k \times k$ diagonal matrix whose ss th element is $x_t(s)$. Let $E(\mathbf{x}_{t+h} | \{s_t\})$ be the $k \times 1$ vector whose st th component is the forecast $E(x_{t+h} | \{s_t = s\})$. Let finally $\mathbf{1}_k$ be the $k \times 1$ unitary vector. Then we have*

$$E(\mathbf{x}_{t+h} | \{s_t\}) = \mathbf{X}_t(\mathbf{\Pi B})^h \mathbf{1}_k.$$

Proof. See in appendix. *Q.E.D.*

The actual dynamics is then obtained by reintroducing the forecast $E(x_{t+h} | \{s_t = s\})$ given in Lemma 8 into (29), that is, for any s ($s = 1, \dots, k$),

$$\sum_{h=1}^H \gamma_h \mathbf{X}_t(\mathbf{\Pi B})^h \mathbf{1}_k + \mathbf{X}_t \mathbf{1}_k + \delta_1 \mathbf{1}_k x_{t-1} = \mathbf{0}_k, \quad (37)$$

where $\mathbf{0}_k$ stands for the $k \times 1$ null vector. The st th equation ($s = 1, \dots, k$) of the system (37) determines implicitly the current state $x_t(s)$ in the event $s_t = s$ as a function

of $\mathbf{\Pi}$, \mathbf{B} , and the relevant past history, here summarized by the previous state x_{t-1} of the system. By definition, in equilibrium, $x_t(s)$ must be equal to $\beta_s x_{t-1}$ in (37) whatever s is ($s = 1, \dots, k$), i.e., $\mathbf{X}_t = \mathbf{B}x_{t-1}$. Under this requirement, (37) rewrites

$$\sum_{h=1}^H \gamma_h \mathbf{B}x_{t-1}(\mathbf{\Pi B})^h \mathbf{1}_k + \mathbf{B}x_{t-1} \mathbf{1}_k + \delta_1 \mathbf{1}_k x_{t-1} = \mathbf{0}_k. \quad (38)$$

Let us now define a sunspot equilibrium on the growth rate as a vector $\boldsymbol{\beta} \equiv (\beta_1, \dots, \beta_k)$ and a Markov matrix $\mathbf{\Pi}$ such that (a) there exist two components β_s and $\beta_{s'}$ of the vector $\boldsymbol{\beta}$ such that $\beta_s \neq \beta_{s'}$ and such that (b) the $k \times 1$ vector $\boldsymbol{\beta} \equiv \mathbf{B}\mathbf{1}_k$ satisfies

$$\sum_{h=0}^H \gamma_h \mathbf{B}(\mathbf{\Pi B})^h \mathbf{1}_k + \delta_1 \mathbf{1}_k = \mathbf{0}_k \quad (39)$$

with the convention that $\gamma_0 \equiv 1$. Equation (39) is obtained from (38) when $x_{t-1} \neq 0$. Condition (a) ensures that the growth rate is changing with sunspot events, while (b) makes the actual growth rate equal to the initial guess β_s in the sunspot event s . If (b) holds true, but (a) is violated, then the solutions to (39) are the $k \times 1$ vectors $\boldsymbol{\beta}$ whose each component β_s is equal to λ_i ($i = 1, \dots, H + 1$).

The mere existence sunspot equilibria associated to a $k \times 1$ vector $\boldsymbol{\beta}$ whose each component β_s ($s = 1, \dots, k$) stands arbitrarily close to some root λ_i would weaken the likelihood that traders succeed in coordinating their forecasts on this root. In view of the fact that any $k \times k$ diagonal matrix $\mathbf{B} = \mathbf{L}_i$ whose ss' th entry is λ_i is a solution to (39) whatever $\mathbf{\Pi}$ is, a $k \times k$ diagonal matrix \mathbf{B} whose ss' th element is arbitrarily close to λ_i will form a sunspot equilibrium on the growth rate associated to $\mathbf{\Pi}$ only if the implicit function theorem breaks at $\mathbf{B} = \mathbf{L}_i$ for the system (39), namely

$$\det \mathbf{D} \left(\sum_{h=0}^H \gamma_h \mathbf{L}_i (\mathbf{\Pi L}_i)^h \mathbf{1}_k + \delta_1 \mathbf{1}_k \right) = 0,$$

where $\mathbf{D}(\bullet)$ is the Jacobian matrix of (\bullet) and $\det \mathbf{D}(\bullet)$ stands for the determinant of this Jacobian matrix. The following result shows that the root of lowest modulus (λ_1) is in fact the only one to be locally immune to sunspots.

Proposition 9 *Let $x_{t-1} \neq 0$ whatever $t \geq 0$ is. Then there do not exist stationary sunspot equilibria on the growth rate $(\boldsymbol{\beta}, \mathbf{\Pi})$ associated with a $k \times 1$ vector $\boldsymbol{\beta}$ whose each component β_s ($s = 1, \dots, k$) stands arbitrarily close to the root of lowest modulus (λ_1). On the contrary there do exist Markov matrices $\mathbf{\Pi}$ such that $(\boldsymbol{\beta}, \mathbf{\Pi})$ is a stationary sunspot equilibria whenever each component β_s ($s = 1, \dots, k$) of $\boldsymbol{\beta}$ stands arbitrarily close to any remaining root λ_i ($i = 2, \dots, H + 1$).*

Proof. See in appendix. *Q.E.D.*

4.3 Learning Growth rates

Assume now that traders try to estimate the law of motion of the level of the state variable restricted to some MSV solution. Their *a priori* belief on the law of motion still fits (3) but β is now estimated by β_t at date t , where β_t is possibly different from some root λ_i of the characteristic polynomial of (30). Therefore their time t forecast $E(x_{t+h} | I_t)$, $h \geq 1$, obtained by leading (3) h times forward, is equal to $\beta_t^h x_t$. The actual dynamics then results from reinserting this forecast into the temporary equilibrium map (29),

$$x_t = -(\delta_1 / (1 + \sum_{h=1}^H \gamma_h \beta_t^h)) x_{t-1}. \quad (40)$$

According to the myopic learning rule, traders must revise at time $(t+1)$ their time t estimate β_t by choosing as new estimate β_{t+1} the actual growth rate at period t , such as given by (40)⁸⁸

$$\beta_{t+1} = -(\delta_1 / (1 + \sum_{h=1}^H \gamma_h \beta_t^h)). \quad (41)$$

The $(H+1)$ roots λ_i are the fixed points of (41). The dynamics with learning (41) arbitrarily close to λ_i is governed by the H roots of the characteristic polynomial associated with the H -dimensional difference equation obtained by linearizing (41) at point $\beta_t = \lambda_i$. As usual, a fixed point λ_i is locally stable under learning if and only if all these H roots have modulus less than 1. The following result provides a condition under which λ_i is locally stable in the dynamics (41).

Proposition 10 *A root λ_i ($i = 1, \dots, H+1$) is locally stable in the dynamics with learning (41) if its modulus $|\lambda_i|$, which measures the speed of convergence of the level of the state variable toward its steady state value along the corresponding MSV solution, is low enough.*

Proof. See in appendix. *Q.E.D.*

In general, unlike the previous sections, the local stability under learning of a solution is not equivalent to its local determinacy. Nevertheless, if the level of the state variable converges rapidly enough toward its steady state along the MSV solution corresponding to the root λ_1 , which is satisfied if, e.g., the influence of past onto the economic system is low enough (δ_1 is close enough to 0), then agents will locally learn this solution, thus making local stability under learning consistent with local determinacy and local immunity to sunspots. However, of course, it is still possible

⁸ Here again note that the algorithm (41) is identical to the iterative version of the expectational stability criterion.

that several MSV solutions to (29) are locally stable in the learning dynamics (41),⁹ but, in this case, the set of stable solutions necessarily includes the MSV solution corresponding λ_1 .

5 Concluding Comments

Rational expectations models typically feature a multiplicity of equilibrium paths. The aim of this paper was to describe how to apply three selection devices, respectively based on local determinacy, local immunity to sunspots and local stability under learning, to MSV solutions of general linear models. The results are clear: All these devices are in general equivalent, and, furthermore, recommend to select a unique MSV solution. This solution is actually the one that is usually believed to be of practical relevance in macroeconomics (in particular, it is the saddle stable path in the saddle point configuration, i.e., the only stable solution in this configuration). There are some open questions that may deserve to be the subject of further work. First formal extensions include the case of the general linear model with an arbitrary number of leads and lags, where the state variable is multidimensional. In view of the results presented in this paper, one can expect most of the dynamic equivalence principle to be saved in such frameworks. Another direction consists in applying the techniques of the paper to bubble solutions, along which the current state of the system depends on more lagged variables than the number of initial conditions to the economic system. In this case the actual equilibrium trajectory is characterized by not only a vector of stationary extended growth rate, but also by some arbitrary forecasts about the level of the state variable in the initial periods. Intuitively, coordination on this type of solutions is much more demanding than coordination on MSV solutions, but there are examples in the literature where bubble solutions are locally stable under learning (see, e.g., Evans and Honkapohja 1994). Therefore, it would be interesting to know whether these solutions can be also locally determinate or immune to sunspots. If this is not the case, as one may conjecture, the relevance of the MSV solution whose role has been recurrently emphasized throughout this paper would still raise in practical analysis.

⁹ Evans and Honkapohja 1994 have actually shown that the class of models (29) with $H = 2$ and $L = 1$ may have two distinct MSV solutions that are locally stable under least squares learning.

A Proof of Proposition 7

The characteristic polynomial Q of the H th order linear recursive equation (36) is

$$Q(\mu) = \sum_{h=0}^H \sum_{j=h}^H \gamma_j \lambda_i^j \mu^h. \quad (42)$$

The root λ_i is locally determinate in the dynamics (35) if and only if all the roots of Q have modulus greater than 1. In order to find these roots, observe first that

$$Q(\mu) = 0 \Leftrightarrow \frac{Q(\mu)}{\gamma_H(\lambda_i)^H} = \sum_{h=0}^H \sum_{j=h}^H \frac{\gamma_j}{\gamma_H} \lambda_i^{j-H} \mu^h = 0, \quad (43)$$

which is allowed when $\gamma_H \neq 0$ and $\delta \neq 0$ since then $\lambda_i \neq 0$ ($i = 1, \dots, H + 1$). We shall now use the relations between the coefficients of the characteristic polynomial corresponding to (30) and the roots λ_i ($i = 1, \dots, H + 1$), namely (see, e.g., Ramis, Deschamps and Odoux 1974)¹⁰

$$\frac{\gamma_{H-j}}{\gamma_H} = (-1)^j \sigma_j(\mathbf{\Lambda}), \quad \text{for any } j \ (j = 1, \dots, H), \quad (46)$$

where $\mathbf{\Lambda}$ denotes the set of the roots $\{\lambda_1, \dots, \lambda_{H+1}\}$ and $\sigma_j(\mathbf{\Lambda})$ is the j th symmetric polynomial in the set $\mathbf{\Lambda}$, i.e.,

$$\sigma_h(\mathbf{\Lambda}) = \sum_{1 \leq j_1 < \dots < j_h} \lambda_{j_1} \dots \lambda_{j_h}.$$

One can show that

$$\sum_{j=h}^H \frac{\gamma_j}{\gamma_H} (\lambda_i)^{j-H} = (-1)^{H-h} \sigma_{H-h}(\mathbf{\Lambda}(l/i)),$$

¹⁰ Consider the polynomial with real coefficients

$$P(z) = z^n + a_{n-1}z^{n-1} \dots + a_0 \quad (44)$$

whose roots are z_i^* ($i = 1, \dots, n$). By definition, we have

$$P(z) = 0 \Leftrightarrow \prod_{i=1}^n (\lambda - z_i^*) = 0 \Leftrightarrow z^n - \left(\sum_{i=1}^n z_i^* \right) z^{n-1} + \dots + (-1)^n \prod_{i=1}^n (z_i^*) = 0 \quad (45)$$

Identifying (44) and (45) term by term leads to the relations

$$-a_n \left(\sum_{i=1}^n z_i^* \right) = a_{n-1}, \dots, a_n (-1)^n \prod_{i=1}^n (z_i^*) = a_0.$$

If one denotes $\sigma_l(z_1^*, \dots, z_n^*)$ the sum over all the different products of l distinct elements ($l = 1, \dots, n$) of the set $\{z_1^*, \dots, z_n^*\}$, i.e., $\sigma_1(z_1^*, \dots, z_n^*)$ is for instance the sum of the n roots and $\sigma_n(z_1^*, \dots, z_n^*)$ is the product of these n roots. The formula (46) directly comes.

where $\Lambda(l/i)$ is the set of all possible ratio λ_l/λ_i for any $l \neq i$ ($l = 1, \dots, H + 1$) with i given ($i = 1, \dots, H + 1$). It follows that

$$Q(\mu) = 0 \Leftrightarrow \sum_{h=0}^H (-1)^{H-h} \sigma_{H-h}(\Lambda(l/i)) \mu^h = 0.$$

The same argument as in the derivation of (46) shows that the H roots of Q are the H ratios λ_l/λ_i for any $l \neq i$. All these roots have modulus larger than 1 if and only if $i = 1$. *Q.E.D.*

B Proof of Lemma 8

Let $\pi_{ss'}$ denote the probability that $s_{t+1} = s'$ ($s' = 1, \dots, k$) if $s_t = s$ ($s = 1, \dots, k$). This probability is the ss' th element of the $k \times k$ matrix $\mathbf{\Pi}$.

Step 1. Here we prove that (??) is satisfied for $h = 1$. Let for instance $s_t = s$ ($s = 1, \dots, k$). In this case $\pi_{ss'}$ is the probability that $\beta_{t+1} = \beta_{s'}$ ($s' = 1, \dots, k$). Hence the expected growth rate $\bar{\beta}_s$ between t and $t + 1$ is given by

$$\bar{\beta}_s = \sum_{s'=1}^k \pi_{ss'} \beta_{s'}.$$

Since $E(x_{t+1} | \{s_t = s\}) = E(\beta_{t+1} | \{s_t = s\})x_t(s)$, we have also

$$E(x_{t+1} | \{s_t = s\}) = \bar{\beta}_s x_t(s).$$

Observe now that $\bar{\beta}_s$ is the s th component of the $k \times 1$ vector $\mathbf{\Pi B 1}$ and $x_t(s)\bar{\beta}_s$ is consequently the s th component of the $k \times 1$ vector $\mathbf{X}_t \mathbf{\Pi B 1}$, which shows Lemma 8 for $h = 1$.

Step 2. Recall first that $E(x_{t+h} | \{s_t = s\}) = E(\beta_{t+1} \cdots \beta_{t+h} | \{s_t = s\})x_t(s)$. Let our recursion hypothesis be that (??) is satisfied for some given $h \geq 1$, i.e., $E(\beta_{t+1} \cdots \beta_{t+h} | \{s_t = s\})$ is the s th component of $(\mathbf{\Pi B})^h \mathbf{1}$. The result follows by exploiting a simple vector recursion over growth rates expected values. By definition, it is indeed true that

$$E(\beta_{t+1} \cdots \beta_{t+h+1} | \{s_t = s\}) = \sum_{s'=1}^k \pi_{ss'} \beta_{s'} E(\beta_{t+2} \cdots \beta_{t+h+1} | \{s_{t+1} = s'\}). \quad (47)$$

But $E(\beta_{t+2} \cdots \beta_{t+h+1} | \{s_{t+1} = s'\}) = E(\beta_{t+1} \cdots \beta_{t+h} | \{s_t = s'\})$. Thus (47) rewrites:

$$\begin{pmatrix} E(\beta_{t+1} \cdots \beta_{t+h+1} | \{s_t = 1\}) \\ \vdots \\ E(\beta_{t+1} \cdots \beta_{t+h+1} | \{s_t = k\}) \end{pmatrix} = \mathbf{\Pi B} \begin{pmatrix} E(\beta_{t+1} \cdots \beta_{t+h} | \{s_t = 1\}) \\ \vdots \\ E(\beta_{t+1} \cdots \beta_{t+h} | \{s_t = k\}) \end{pmatrix}$$

The result directly comes from both our recursion hypothesis and the fact that $E(x_{t+h+1} | \{s_t = s\}) = E(\beta_{t+1} \cdots \beta_{t+h+1} | \{s_t = s\})x_t(s)$, i.e., $E(\mathbf{x}_{t+h+1} | \{s_t\}) = \mathbf{X}_t (\mathbf{\Pi B})^{h+1} \mathbf{1}$. This proves Lemma 8 for any $h \geq 1$. *Q.E.D.*

C Proof of Proposition 9

Equation (37) describes a system of k equations whose k unknowns are the k growth rates components β_s ($s = 1, \dots, k$) of the $k \times 1$ vector $\boldsymbol{\beta}$. Therefore the Jacobian matrix of this system is of dimension $k \times k$. This matrix is derived by using basic matrix differential calculus. We first define

$$F(\mathbf{B}) = \sum_{h=0}^H \gamma_h \mathbf{B}(\boldsymbol{\Pi}\mathbf{B})^h \mathbf{1} + \delta \mathbf{1}, \quad \text{and} \quad \mathbf{B}(\boldsymbol{\beta}) = \begin{pmatrix} \beta_1 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \beta_k \end{pmatrix}.$$

A sunspot equilibrium is consequently a $k \times 1$ vector $\boldsymbol{\beta}$ associated with a $k \times k$ matrix $\mathbf{B}(\boldsymbol{\beta})$ such that $\boldsymbol{\beta}$ belongs to the kernel of $F(\mathbf{B}(\boldsymbol{\beta}))$. The Jacobian matrix $\mathbf{D}F(\mathbf{B}(\boldsymbol{\beta}))$ at some point $\boldsymbol{\beta}$ of \mathbb{R}^k is equal to $\mathbf{D}F(\mathbf{B})\mathbf{D}\mathbf{B}(\boldsymbol{\beta})$, where $\mathbf{D}F(\mathbf{B})$ and $\mathbf{D}\mathbf{B}(\boldsymbol{\beta})$ are as follows (see 9.4.1 in Magnus and Neudecker 1988)

$$\mathbf{D}F(\mathbf{B}) = \frac{\partial \text{vec}F(\mathbf{B})}{\partial \text{vec}\mathbf{B}}, \quad \text{and} \quad \mathbf{D}\mathbf{B}(\boldsymbol{\beta}) = \frac{\partial \text{vec}\mathbf{B}(\boldsymbol{\beta})}{\partial \text{vec}\boldsymbol{\beta}},$$

where the symbol ∂ stands for the differential and where the vec operator transforms a matrix into a vector by stacking its columns one underneath the other. Of course $\text{vec}F(\mathbf{B})$ is the $k \times 1$ vector $F(\mathbf{B})$. Given that \mathbf{B} is a $k \times k$ matrix, $\text{vec}\mathbf{B}$ is a $k^2 \times 1$ vector. It results that $\mathbf{D}F(\mathbf{B})$ is a $k \times k^2$ matrix. The remaining Jacobian matrix $\mathbf{D}\mathbf{B}(\boldsymbol{\beta})$ is of dimension $k^2 \times k$ since $\text{vec}\boldsymbol{\beta} = \boldsymbol{\beta}$ is a $k \times 1$ vector (and recall that $\text{vec}\mathbf{B}$ is a $k^2 \times 1$ vector). Then, by the chain rule, it follows that $\mathbf{D}F(\mathbf{B}(\boldsymbol{\beta}))$ is a $k \times k$ matrix.

We prove Proposition 9 in two steps. First we provide an expression for $\mathbf{D}F(\mathbf{B}(\boldsymbol{\beta}))$. Second we show that $\det \mathbf{D}F(\mathbf{B}(\boldsymbol{\beta}))$ is always different from zero if β_s ($s = 1, \dots, k$) is equal to λ_1 while there are some stochastic matrices $\boldsymbol{\Pi}$ that make $\mathbf{D}F(\mathbf{B}(\boldsymbol{\beta}))$ singular as soon as β_s ($s = 1, \dots, k$) is equal to λ_i ($i = 2, \dots, H + 1$). Proposition 9 is then a straightforward consequence of the implicit function theorem. These two steps are the subject of two different lemma.

Lemma 11 *The Jacobian matrix $\mathbf{D}F(\mathbf{B}(\boldsymbol{\beta}))$ is given by*

$$\mathbf{D}F(\mathbf{B}(\boldsymbol{\beta})) = \sum_{h=0}^H \gamma_h \lambda_i^h \sum_{j=0}^h \boldsymbol{\Pi}^j, \quad \text{at point } \boldsymbol{\beta} = (\lambda_i, \dots, \lambda_i)'$$

Proof. The map F has the following differential

$$\partial F(\mathbf{B}) = \sum_{h=0}^H \gamma_h \partial \left[\mathbf{B}(\boldsymbol{\Pi}\mathbf{B})^h \right] \mathbf{1}, \quad (48)$$

where $(\mathbf{\Pi B})^h$ stands for the product $(\mathbf{\Pi B}) \cdots (\mathbf{\Pi B})$ h times. A mild adaptation of results in table 7 (Magnus and Neudecker 1988, Ch. 9) allows us to get

$$\partial [\mathbf{B\Pi B} \cdots \mathbf{\Pi B}] = \sum_{j=0}^h (\mathbf{B\Pi})^j \partial \mathbf{B} (\mathbf{\Pi B})^{h-j}.$$

As a result (48) rewrites

$$\begin{aligned} \partial F(\mathbf{B}) &= \sum_{h=0}^H \gamma_h \sum_{j=0}^h (\mathbf{B\Pi})^j \partial \mathbf{B} (\mathbf{\Pi B})^{h-j} \mathbf{1} \\ \Rightarrow \text{vec} \partial F(\mathbf{B}) &= \partial \text{vec} F(\mathbf{B}) = \sum_{h=0}^H \gamma_h \sum_{j=0}^h \text{vec} [(\mathbf{B\Pi})^j \partial \mathbf{B} (\mathbf{\Pi B})^{h-j} \mathbf{1}]. \end{aligned}$$

Now it follows from theorem 2.2 in Magnus and Neudecker 1988 that

$$\partial \text{vec} F(\mathbf{B}) = \sum_{h=0}^H \gamma_h \sum_{j=0}^h [((\mathbf{\Pi B})^{h-j} \mathbf{1})' \otimes (\mathbf{B\Pi})^j] \partial \text{vec} \mathbf{B},$$

where the symbol $'$ represents the matrix transpose and the symbol \otimes stands for the Kronecker product. Let us take now $\mathbf{B} = \mathbf{\Lambda}_i = \lambda_i \mathbf{I}_k$ where λ_i is some root of the characteristic equation associated with (30). Then one obtains

$$\partial \text{vec} F(\mathbf{\Lambda}_i) = \sum_{h=0}^H \gamma_h \lambda_i^h \sum_{j=0}^h [(\mathbf{1}' (\mathbf{\Pi}')^{h-j}) \otimes \mathbf{\Pi}^j] \partial \text{vec} \mathbf{\Lambda}_i.$$

Observe that the inner product between $\mathbf{1}'$ and the s th column of $(\mathbf{\Pi}')^{h-j}$ is merely equal to the sum over all the components of the s th row of $\mathbf{\Pi}^{h-j}$, that is equal to 1 by definition of a Markov matrix, i.e., $\mathbf{1}' (\mathbf{\Pi}')^{h-j} = \mathbf{1}'$. Hence

$$\partial \text{vec} F(\mathbf{\Lambda}_i) = \sum_{h=0}^H \gamma_h \lambda_i^h \sum_{j=0}^h \overbrace{(\mathbf{\Pi}^j \cdots \mathbf{\Pi}^j)}^k \partial \text{vec} \mathbf{\Lambda}_i, \quad (49)$$

where $(\mathbf{\Pi}^j \cdots \mathbf{\Pi}^j)$ is the $k \times k^2$ matrix whose ss' th element ($s, s' = 1, \dots, k$), i.e., the ss' th element of the $k \times k$ matrix $\mathbf{\Pi}^j$, is the same as its $s(s' + k \bmod n)$ th element ($n = 1, \dots, k$). Given the way the Jacobian matrix of F at point $\mathbf{\Lambda}_i$ is defined, it directly follows from (49) that

$$\mathbf{D}F(\mathbf{\Lambda}) = \sum_{h=0}^H \gamma_h \lambda^h \sum_{j=0}^h \overbrace{(\mathbf{\Pi}^j \cdots \mathbf{\Pi}^j)}^k.$$

We now come to calculate the remaining Jacobian matrix $\mathbf{DB}(\boldsymbol{\beta})$. Observe that

$$\partial \text{vec} \mathbf{B}(\boldsymbol{\beta}) = \begin{pmatrix} \partial \beta_1 \\ 0 \\ \vdots \\ 0 \\ \partial \beta_2 \\ 0 \\ \vdots \\ \partial \beta_k \end{pmatrix} = \begin{pmatrix} 1 & 0 & \cdots & \cdots & 0 \\ 0 & \vdots & & & \vdots \\ \vdots & \vdots & & & \vdots \\ 0 & \vdots & & & \vdots \\ 0 & 1 & 0 & \cdots & 0 \\ 0 & 0 & 0 & \cdots & 0 \\ \vdots & \vdots & & & \vdots \\ 0 & 0 & \cdots & 0 & 1 \end{pmatrix} \begin{pmatrix} \partial \beta_1 \\ \partial \beta_2 \\ \vdots \\ \vdots \\ \vdots \\ \vdots \\ \vdots \\ \partial \beta_k \end{pmatrix} \equiv \boldsymbol{\Phi} \partial \text{vec} \boldsymbol{\beta}.$$

By definition it follows that $\mathbf{DB}(\boldsymbol{\beta}) = \boldsymbol{\Phi}$, where $\boldsymbol{\Phi}$ is a $k^2 \times k$ matrix. Recall now that the chain rule applies and notice that the ss' th element of the $k \times k$ matrix $(\boldsymbol{\Pi}^j \cdots \boldsymbol{\Pi}^j) \boldsymbol{\Phi}$ is the ss' th element of the $k \times k$ matrix $\boldsymbol{\Pi}^j$. Therefore we have, for $\boldsymbol{\beta} = (\lambda_i, \dots, \lambda_i)'$,

$$\mathbf{DF}(\mathbf{B}(\boldsymbol{\beta})) = \sum_{h=0}^H \gamma_h \lambda_i^h \sum_{j=0}^h (\boldsymbol{\Pi}^j \boldsymbol{\Pi}^j) \boldsymbol{\Phi} = \sum_{h=0}^H \gamma_h \lambda_i^h \sum_{j=0}^h \boldsymbol{\Pi}^j.$$

and Lemma 11 follows. *Q.E.D.*

Lemma 12 *Let π_s ($s = 1, \dots, k$) be an eigenvalue of the Markov matrix $\boldsymbol{\Pi}$. Let \mathbf{l}_i be the $k \times 1$ vector whose all the components are equal to a given root λ_i ($i = 1, \dots, H+1$). Assume that $\lambda_i \neq 0$. Then the $k \times k$ Jacobian matrix $\mathbf{DF}(\boldsymbol{\Lambda}(\bullet))$ is singular at point \mathbf{l}_i if and only there exists an eigenvalue π_s of $\boldsymbol{\Pi}$ such that*

$$\prod_{\substack{j=1 \\ j \neq i}}^{H+1} \left(\pi_s - \frac{\lambda_j}{\lambda_i} \right) = 0.$$

Proof. Let ϕ_s ($s = 1, \dots, k$) be an eigenvalue of $\mathbf{DF}(\boldsymbol{\Lambda}(\bullet))$ at point \mathbf{l}_i and recall that the matrix $\mathbf{DF}(\boldsymbol{\Lambda}(\bullet))$ is singular at point \mathbf{l}_i if and only if its determinant $\det \mathbf{DF}(\boldsymbol{\Lambda}(\bullet))$ equals 0 at this point. Let now $\boldsymbol{\pi}$ be an eigenvector of $\boldsymbol{\Pi}$ associated with the eigenvalue π . Observe that

$$\mathbf{DF}(\boldsymbol{\Lambda}(\mathbf{l}_i)) \boldsymbol{\pi} = \sum_{h=0}^H \gamma_h \lambda_i^h \sum_{j=0}^h \boldsymbol{\Pi}^j \boldsymbol{\pi} = \left(\sum_{h=0}^H \gamma_h \lambda_i^h \sum_{j=0}^h \boldsymbol{\pi}^j \right) \boldsymbol{\pi},$$

which shows that the eigenvalues of $\mathbf{DF}(\boldsymbol{\Lambda}(\bullet))$ at point \mathbf{l}_i are of the following form

$$\phi = \sum_{h=0}^H \gamma_h \lambda_i^h \sum_{j=0}^h \pi^j.$$

Of course $\det \mathbf{D}F(\mathbf{\Lambda}(\bullet))$ equals 0 at point \mathbf{l}_i if and only if one of the eigenvalues ϕ of $\mathbf{D}F(\mathbf{\Lambda}(\mathbf{l}_i))$ is equal to 0, namely

$$\sum_{h=0}^H \gamma_h \lambda_i^h \sum_{j=0}^h \pi^j = 0 \Leftrightarrow \sum_{h=0}^H \pi^h \sum_{j=h}^H \gamma_h \lambda_i^h = 0. \quad (50)$$

Let us now divide the right member of (50) by $\gamma_H \lambda_i^H$ (which is assumed to differ from 0). One gets

$$\sum_{h=0}^H \pi^h \sum_{j=h}^H \frac{\gamma_h \lambda_i^h}{\gamma_H \lambda_i^H} = 0 \Leftrightarrow \prod_{\substack{j=1 \\ j \neq i}}^{H+1} \left(\pi - \frac{\lambda_j}{\lambda_i} \right) = 0.$$

where the last equality comes from Proposition 7 (see equation 43 in particular), which proves Lemma 12. *Q.E.D.*

Given that the real part of any eigenvalue of a Markov matrix has modulus less than 1, i.e., $|\pi_s| < 1$, and given that the roots λ_i are labelled in the order of increasing modulus, the ratio $|\lambda_j/\lambda_i|$ is strictly greater than 1 for $j \neq 1$ ($j = 1, \dots, H+1$). Proposition 9 follows from the implicit function theorem since $\lambda_1 \neq 0$ under the assumption that $\delta_1 \neq 0$, and standard local bifurcation theory (see Chiappori, Geoffard and Guesnerie 1992 for a general argument). *Q.E.D.*

D Proof of Proposition 10

The dynamics with learning (41) arbitrarily close to the root λ_i ($i = 1, \dots, H+1$) is obtained by linearizing (41) at point $\beta_{t+1} = \beta_t = \lambda_i$. This dynamics is governed by the ratio

$$\frac{d\beta_{t+1}}{d\beta_t} = \delta_1 \frac{\sum_{h=1}^H h \gamma_h \lambda_i^{h-1}}{(1 + \sum_{h=1}^H \gamma_h \lambda_i^h)^2}.$$

The root λ_i is locally stable under learning if and only if this ratio is less than 1 in modulus. This is the case when λ_i is close enough to 0, since then $|d\beta_{t+1}/d\beta_t|$ also is close to 0. *Q.E.D.*

References

- [1] Azariadis, Costas (1981) Self-fulfilling prophecies. *Journal of Economic Theory* 25, 380-96.
- [2] Azariadis, Costas and Roger Guesnerie (1982) Prophéties créatrices et persistance des théories. *Revue Economique* 33, 787-806.
- [3] Blanchard, Olivier J. and Charles Kahn (1981) The solution of linear difference models under rational expectations. *Econometrica* 48, 1305-1311.
- [4] Chiappori, Pierre-André, Pierre-Yves Geoffard and Roger Guesnerie (1992) Sunspot fluctuations around a steady state: the case of multidimensional one-step forward looking models. *Econometrica* 60, 1097-1126.
- [5] DeCanio, Stephen (1979) Rational expectations and learning from experience. *Quarterly Journal of Economics* 93, 47-58.
- [6] Desgranges, Gabriel and Stéphane Gauthier (in press) On the uniqueness of the bubble-free solution in linear rational expectations models. *Macroeconomic Dynamics*.
- [7] Evans, George (1985) Expectational stability and the multiple equilibria problem in linear rational expectations models. *Quarterly Journal of Economics* 100, 147-157.
- [8] Evans, George (1989) The fragility of sunspots and bubbles. *Journal of Monetary Economics* 23, 297-317.
- [9] Evans, George and Roger Guesnerie (1993) Rationalizability, strong rationality, and expectational stability. *Games and Economic Behavior* 5, 632-646.
- [10] Evans, George and Seppo Honkapohja (1994) Learning, convergence, and stability with multiple rational expectations equilibria. *European Economic Review* 38, 1071-1098.
- [11] Evans, George and Seppo Honkapohja (1999) Learning dynamics. In John B. Taylor and Michael Woodford (ed.), *Handbook of Macroeconomics*, vol. 1, pp. 449-542. Elsevier Science.
- [12] Farmer, Roger and Michael Woodford (1997) Self-fulfilling prophecies and the business cycle. *Macroeconomic Dynamics* 1, 740-769.
- [13] Gantmacher, Felix R. (1966), *Théorie des Matrices*, Editions Jacques Gabay, Paris: Dunod.

- [14] Gauthier, Stéphane (in press) Determinacy and stability under learning of rational expectations equilibria. *Journal of Economic Theory*.
- [15] Grandmont, Jean-Michel (1998) Expectations formation and stability of large socioeconomic systems. *Econometrica* 66, 741-783.
- [16] Grandmont, Jean-Michel and Guy Laroque (1986) Stability of cycles and expectations. *Journal of Economic Theory* 40, 138-151.
- [17] Grandmont, Jean-Michel and Guy Laroque (1991) Economic dynamics with learning: some instability examples. In William Barnett (ed.), *Equilibrium Theory and Applications: Proceedings of the Sixth International Symposium in Economic Theory and Econometrics*. Cambridge U.K., Cambridge University Press.
- [18] Guesnerie, Roger (1993) Theoretical tests of the rational expectations hypothesis in economic dynamic models. *Journal of Economic Dynamics and Control* 17, 847-864.
- [19] Guesnerie, Roger and Michael Woodford (1991) Stability of cycles under adaptive learning rules. In William Barnett (ed.), *Proceedings of the Sixth International Symposium in Economic Theory and Econometrics*. Cambridge U.K.: Cambridge University Press.
- [20] Kehoe, Timothy J. and David Levine (1985) Comparative Statics and Perfect Foresight in Infinite Horizon, *Econometrica* 53, 433-454.
- [21] McCallum, Bennett T. (1983) On non-uniqueness in rational expectations models: an attempt at perspective. *Journal of Monetary Economics* 11, 139-168.
- [22] McCallum, Bennett (1999) Role of the minimal state variable criterion in rational expectations models. NBER Working Paper 7087.
- [23] Magnus, Jan R. and Heinz Neudecker (1988) *Matrix differential calculus with applications to statistics and econometrics*. Wiley series in probability and mathematical statistics.
- [24] Marcet, Albert and Thomas Sargent (1989) Convergence of least-squares learning mechanisms in self-referential stochastic models. *Journal of Economic Theory* 48, 337-368.
- [25] Mignotte, Maurice (1989) *Mathématiques pour le calcul formel*. Paris: Presses Universitaires de France.
- [26] Ramis, Edmond, Claude Deschamps and Jacques Odoux (1974) *Cours de mathématiques spéciales*, Tome 1. Paris: Masson.

- [27] Taylor, John B. (1977) Conditions for Unique Solutions in Stochastic Macroeconomic Models with Rational Expectations. *Econometrica* 45, 1377-1385.
- [28] Woodford, Michael (1984) Indeterminacy of equilibrium in the overlapping generations model: a survey. Mimeo: Columbia University.