



HAL
open science

The cone of supermodular games on finite distributive lattices

Michel Grabisch, Tomáš Kroupa

► **To cite this version:**

Michel Grabisch, Tomáš Kroupa. The cone of supermodular games on finite distributive lattices. *Discrete Applied Mathematics*, 2019, 260, pp.144-154. 10.1016/j.dam.2019.01.024 . halshs-02381019v2

HAL Id: halshs-02381019

<https://shs.hal.science/halshs-02381019v2>

Submitted on 8 May 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

The cone of supermodular games on finite distributive lattices

Michel Grabisch^a, Tomáš Kroupa^{b,*}

^aParis School of Economics, University of Paris I, Paris, France

^bThe Czech Academy of Sciences, Institute of Information Theory and Automation, Prague, Czech Republic

Abstract

In this article we study supermodular functions on finite distributive lattices. Relaxing the assumption that the domain is a powerset of a finite set, we focus on geometrical properties of the polyhedral cone of such functions. Specifically, we generalize the criterion for extremality and study the face lattice of the supermodular cone. An explicit description of facets by the corresponding tight linear inequalities is provided.

Keywords: supermodular/submodular function, core, coalitional game, polyhedral cone

1. Introduction

Supermodular functions, and their duals, submodular functions, play a central rôle in many fields of discrete mathematics, most notably combinatorial optimization (rank function of polymatroids: see, e.g., the monograph of Fujishige [8]), game theory (characteristic function of transferable utility games: see, e.g., Peleg and Sudhölter [14]), decision theory (capacity, Choquet expected utility [18]), lattice theory, etc.

Up to duality, all above examples fall into the category of *supermodular games*, that is, supermodular functions vanishing at the empty set. They form a polyhedral cone, whose facets have been found by Kuipers et al. [13]. In his 1971 seminal paper, Shapley [19] gave the 37 extreme supermodular games for $n = 4$ players, and noted that for larger values of n , little can be said. Later, Rosenmüller and Weidner [17] found all extreme supermodular functions by representing each such function as a maximum over shifted additive games. Recently, Studený and Kroupa [21] revisited the problem and provided another characterization of extremality, in a sense dual to the result of Rosenmüller and Weidner, but easier to use.

The aim of this paper is to (re)establish in a more general framework and in a simpler way the above results (together with new ones) describing the cone of supermodular games, taking advantage of classical results on polyhedra. We consider games defined on a finite distributive lattice \mathcal{L} , generated by a partial order \preceq on the set of players N . The poset induces some relation between the players, which can be interpreted in various

*Corresponding author

Email addresses: michel.grabisch@univ-paris1.fr (Michel Grabisch), kroupa@utia.cas.cz (Tomáš Kroupa)

ways: precedence constraints (Faigle and Kern [6]), hierarchy (Grabisch and Xie [11]), or permission structure (van den Brink and Gilles [4]). Feasible coalitions of players, i.e., those for which the game is defined, are down-sets on (N, \preceq) , and they form a distributive lattice \mathcal{L} . By Birkhoff's theorem, every finite distributive lattice is of this form. The standard case $\mathcal{L} = 2^N$ is recovered when the poset (N, \preceq) is flat, i.e., when all players are incomparable (no order relation between the players).

A large amount of research has been done concerning games on distributive lattices, as well as on other ordered structures (see a survey in [9]). Most of them are related to the solution concepts such as Shapley value or the core. However, up to our knowledge, there is no systematic study on the geometric properties of the cone of supermodular games defined on distributive lattices. Note that it is very natural to take a distributive lattice as a domain of a supermodular function since supermodular inequalities involve only the lattice joins and meets. The present paper addresses precisely this point. In the same time we generalize and prove results about extreme rays and facets in a more concise way.

Section 2 collects background on distributive lattices. Coalitional games are introduced in Section 3, in particular 0-normalized and supermodular games. Section 4 contains basic facts about 0-normalized supermodular games and the cone thereof. The extreme rays are characterized in Section 5. Basically, a supermodular game generates an extreme ray if and only if a certain system of linear equalities has for a solution those vectors which are proportional to the marginal vectors of the game. Section 6 describes the facial structure of the cone by a certain collection of finite lattices, namely the tight sets associated with compatible permutations of the poset (N, \preceq) . The facets of the cone of supermodular games are characterized in Section 7.

2. Finite distributive lattices

In this section we introduce basic notions and results about Birkhoff duality between finite distributive lattices and finite posets. The reader is referred to [20, Chapter 3] for all the unexplained notions concerning lattices and partially ordered sets (posets).

Let \mathcal{L} be a finite distributive lattice whose join and meet are denoted by \vee and \wedge , respectively. A partial order \leq on \mathcal{L} is defined by $a \leq b$ if $a \vee b = b$, for all $a, b \in \mathcal{L}$. Since \mathcal{L} is finite there exists a top element \top and a bottom element \perp in \mathcal{L} . We always assume that \mathcal{L} is non-trivial in sense that $\top \neq \perp$. An element $a \in \mathcal{L}$ is called *join-irreducible* if $a \neq \perp$ and the identity $a = b \vee c$ holding for some $b, c \in \mathcal{L}$ implies $a = b$ or $a = c$. In particular, $a \in \mathcal{L}$ with $a \neq \perp$ is an *atom* if the condition $b \leq a$ for all $b \in \mathcal{L}$ implies $b = \perp$ or $b = a$. The join-irreducible elements of a Boolean lattice are precisely its atoms. For any $a, b \in \mathcal{L}$ such that $a \leq b$, we define an *order interval*

$$[a, b] := \{c \in \mathcal{L} \mid a \leq c \leq b\}.$$

An element $a \in \mathcal{L}$ is join-irreducible if, and only if, there is a unique $a^- \in \mathcal{L}$ such that $a^- \leq a$, $a^- \neq a$, and $[a^-, a] = \{a^-, a\}$. The set of all join-irreducible elements of \mathcal{L} is denoted by $\mathcal{J}(\mathcal{L})$ and it is always endowed with the partial order \leq of \mathcal{L} restricted to $\mathcal{J}(\mathcal{L})$. Thus, $(\mathcal{J}(\mathcal{L}), \leq)$ becomes a nonempty finite poset.

Let $N \neq \emptyset$ be a finite set and \preceq be a partial order on N . A *down-set* in (N, \preceq) is a subset $A \subseteq N$ such that if $i \in A$ and $j \preceq i$ for $j \in N$, then $j \in A$. For any $i \in N$, we denote

$$\downarrow i := \{j \in N \mid j \preceq i\} \quad \text{and} \quad \Downarrow i := \downarrow i \setminus \{i\}.$$




Figure 2.1: The poset (N, \preceq) with the corresponding lattice of down-sets $\mathcal{D}(N, \preceq)$

Both $\downarrow i$ and $\downarrow j$ are down-sets in (N, \preceq) . A down-set A is called *principal* if there exists some $i \in N$ such that $A = \downarrow i$. By $\mathcal{D}(N, \preceq)$ we denote the set of all down-sets in (N, \preceq) . It is easy to see that $\mathcal{D}(N, \preceq)$ is closed under the set-theoretic union \cup and intersection \cap . Thus, $\mathcal{D}(N, \preceq)$ is a finite distributive lattice whose order is the inclusion \subseteq between sets, and whose top and bottom element is N and \emptyset , respectively. The lattice $\mathcal{D}(N, \preceq)$ is the most general example of a finite distributive lattice by the following classical result.

Birkhoff's representation theorem. *Let \mathcal{L} be a finite distributive lattice. Then the mapping*

$$F_{\mathcal{L}}: \mathcal{L} \rightarrow \mathcal{D}(\mathcal{J}(\mathcal{L}), \leq)$$

defined by $F_{\mathcal{L}}(a) := \{b \in \mathcal{J}(\mathcal{L}) \mid b \leq a\}$ is a lattice isomorphism.

The converse part of duality explains what are join-irreducible elements in the lattice of down-sets $\mathcal{D}(N, \preceq)$.

Proposition 2.1. *Let (N, \preceq) be a finite poset. Then the mapping*

$$\downarrow: (N, \preceq) \rightarrow \mathcal{J}(\mathcal{D}(N, \preceq))$$

sending every $i \in N$ to the principal down-set $\downarrow i$ is an order isomorphism.

Example 2.1. Let $N := \{1, 2, 3, 4\}$ be equipped with the partial order \preceq captured by the Hasse diagram in Figure 2.1 on the left. On the right-hand side we depict the lattice of down-sets $\mathcal{D}(N, \preceq)$. There are four join-irreducible elements in $\mathcal{D}(N, \preceq)$, namely $\{2\}$, $\{3\}$, $\{4\}$, and $\{1, 2, 3\}$.

3. Coalitional games on finite distributive lattices

We use the standard terminology of cooperative game theory; see [14]. The *player set* is defined to be $N := \{1, \dots, n\}$, for some integer $n \geq 1$. Any subset of N is called a *coalition*. We allow for a situation in which players $i, j \in N$ are compared using a

partial order \preceq on N . Hence, (N, \preceq) is assumed to be a finite poset. Birkhoff duality (see Section 2) entails that the partial order \preceq on N restricts the formation of coalitions $A \subseteq N$, provided that the coalition structure is modeled by the lattice of down-sets in (N, \preceq) .

Convention. *Throughout the paper we will always assume that the set of all possible coalitions in (N, \preceq) is the lattice of down-sets $\mathcal{D}(N, \preceq)$. We use the abbreviations*

$$\mathcal{L} := \mathcal{D}(N, \preceq) \quad \text{and} \quad \mathcal{J} := \mathcal{J}(\mathcal{D}(N, \preceq)).$$

From now on, all possible coalitions are assumed to be precisely the sets belonging to a fixed lattice \mathcal{L} and \mathcal{J} denotes its subset of all join-irreducible elements. Coalitional games are modeled as real functions v on the set \mathcal{L} of feasible coalitions A , where the real value $v(A)$ indicates the amount of utility resulting from the joint cooperation of players in the coalition A .

Definition 3.1. A function $v: \mathcal{L} \rightarrow \mathbb{R}$ satisfying $v(\emptyset) = 0$ is a (*coalitional*) *game*. A game v is called

- *supermodular* if $v(A \cup B) + v(A \cap B) \geq v(A) + v(B)$,
- *modular* if $v(A \cup B) + v(A \cap B) = v(A) + v(B)$,
- *monotone* if $v(A) \leq v(B)$ whenever $A \subseteq B$,
- *nonnegative* if $v(A) \geq 0$,

for all $A, B \in \mathcal{L}$.

Let $G(\mathcal{L})$ be the set of all games on \mathcal{L} . We consider these subsets of $G(\mathcal{L})$:

$$\begin{aligned} G_S(\mathcal{L}) &:= \{v \in G(\mathcal{L}) \mid v \text{ is supermodular}\}, \\ G_M(\mathcal{L}) &:= \{v \in G(\mathcal{L}) \mid v \text{ is modular}\}. \end{aligned}$$

A modular game is also called a *valuation* (over \mathbb{R}) in literature; see [2, 20]. Note that $G(\mathcal{L})$ is a real vector space isomorphic to $\mathbb{R}^{\mathcal{L} \setminus \{\emptyset\}}$ and therefore $\dim G(\mathcal{L}) = |\mathcal{L}| - 1$. One of the bases in $G(\mathcal{L})$ is found very easily. For each nonempty $A \in \mathcal{L}$, the *unanimity game* u_A is defined by

$$u_A(B) := \begin{cases} 1 & A \subseteq B, \\ 0 & \text{otherwise,} \end{cases} \quad \text{for all } B \in \mathcal{L}.$$

Then $\{u_A \mid \emptyset \neq A \in \mathcal{L}\}$ forms a basis in $G(\mathcal{L})$. The coordinates of any game $v \in G(\mathcal{L})$ with respect to this basis are calculated using the Möbius inversion formula [15]. Specifically, the *Möbius function* of \mathcal{L} is the function $\mu_{\mathcal{L}}: \mathcal{L}^2 \rightarrow \mathbb{R}$ given recursively as

$$\mu_{\mathcal{L}}(X, Y) := \begin{cases} 1 & X = Y, \\ - \sum_{X \subseteq Z \subset Y} \mu_{\mathcal{L}}(X, Z) & X \subset Y, \\ 0 & \text{otherwise,} \end{cases} \quad \text{for all } X, Y \in \mathcal{L}.$$

The *Möbius transform* of $v \in G(\mathcal{L})$ is the game $\hat{v} \in G(\mathcal{L})$ defined by

$$\hat{v}(B) := \sum_{C \subseteq B} v(C) \cdot \mu_{\mathcal{L}}(C, B), \quad B \in \mathcal{L}.$$

Lemma 3.1. For any $v \in G(\mathcal{L})$, we have

$$\hat{v}(B) = \sum_{C \subseteq B} v(C) \cdot (-1)^{|B \setminus C|}, \quad B \in \mathcal{L},$$

where the sum above is over all $C \in \mathcal{L}$ such that $[C, B]$ is a Boolean sublattice of \mathcal{L} .

Proof. It suffices to apply the observation from [20, Example 3.9.6]. Specifically, since the lattice \mathcal{L} is finite and distributive, the formula for Möbius function $\mu_{\mathcal{L}}$ simplifies as

$$\mu_{\mathcal{L}}(X, Y) = \begin{cases} (-1)^{|Y \setminus X|} & \text{if } [X, Y] \text{ is a Boolean lattice,} \\ 0 & \text{otherwise,} \end{cases}$$

for every $X, Y \in \mathcal{L}$ with $X \subseteq Y$. □

Thus, any $v \in G(\mathcal{L})$ can be expressed as a linear combination $v = \sum_{\emptyset \neq A \in \mathcal{L}} \hat{v}(A) \cdot u_A$, which gives

$$v(A) = \sum_{B \subseteq A} \hat{v}(B), \quad A \in \mathcal{L}. \quad (3.1)$$

The set of valuations (modular games) $G_M(\mathcal{L})$ is a vector subspace of $G(\mathcal{L})$. By Rota's lemma [16] any valuation on \mathcal{L} is uniquely determined by its restriction to the set of join-irreducible elements \mathcal{J} . It follows that the dimension of linear space $G_M(\mathcal{L})$ equals $|\mathcal{J}| = n$. This means that the polyhedral cone of supermodular games $G_S(\mathcal{L})$ is not pointed as it includes the non-trivial linear space $G_M(\mathcal{L})$. However, we can always consider the elements of $G_S(\mathcal{L})$ modulo $G_M(\mathcal{L})$. To this end we introduce the following notion.

Definition 3.2. A game $v \in G(\mathcal{L})$ is said to be *0-normalized* if

$$\hat{v}(A) = 0, \quad \text{for all } A \in \mathcal{J}.$$

Let $G^*(\mathcal{L})$ be the set of all 0-normalized games on \mathcal{L} .

Note that the notion of 0-normalized game on a distributive lattice \mathcal{L} coincides with the usual concept of 0-normalized game in cooperative game theory (see [14, Definition 2.1.13]). Indeed, when \mathcal{L} is the Boolean lattice 2^N of all subsets of N , then the only join-irreducible elements in 2^N are exactly the atoms in 2^N , that is, $\mathcal{J} = \{\{i\} \mid i \in N\}$. If v is 0-normalized in sense of Definition 3.2, from (3.1) we get $v(\{i\}) = \hat{v}(\{i\}) = 0$ for all $i \in N$, which is exactly the definition of 0-normalized coalitional game on 2^N .

Lemma 3.2. A game $v \in G(\mathcal{L})$ is 0-normalized if and only if $v(A) = v(A^-)$ for all $A \in \mathcal{J}$, where A^- is the unique element covered by A .

Proof. Let $A \in \mathcal{J}$. Then, for any $B \in \mathcal{L}$ with $B \subseteq A$, the order interval $[B, A]$ is a Boolean sublattice of \mathcal{L} if, and only if, either $B = A$ or $B = A^-$. Hence, Lemma 3.1 yields $\hat{v}(A) = v(A) - v(A^-)$. □

For any $v \in G(\mathcal{L})$, put

$$v^* := v - \sum_{B \in \mathcal{J}} \hat{v}(B) \cdot u_B.$$

It is easy to see that $G^*(\mathcal{L}) = \{v^* \mid v \in G(\mathcal{L})\}$. We claim that, for any $v \in G(\mathcal{L})$, there exist uniquely determined $w \in G^*(\mathcal{L})$ and $m \in G_M(\mathcal{L})$ such that

$$v = w + m. \quad (3.2)$$

Indeed, it suffices to define $w := v^*$, $m := \sum_{B \in \mathcal{J}} \hat{v}(B) \cdot u_B$, and observe that $m \in G_M(\mathcal{L})$. Let

$$G_S^*(\mathcal{L}) := G_S(\mathcal{L}) \cap G^*(\mathcal{L}).$$

Then $G_S^*(\mathcal{L}) = \{v^* \mid v \in G_S(\mathcal{L})\}$ and Lemma 3.2 says that $G_S^*(\mathcal{L})$ contains exactly those supermodular games satisfying $v(A) = v(A^-)$, for all $A \in \mathcal{J}$. Moreover, the convex cone $G_S^*(\mathcal{L})$ is pointed and polyhedral.

Lemma 3.3. *Every game $v \in G_S^*(\mathcal{L})$ is monotone and nonnegative.*

Proof. Since a monotone game is necessarily nonnegative, it suffices to check monotonicity. We only need to prove that for all $A, B \in \mathcal{L}$ satisfying $B \subseteq A$ and $|B| = |A| - 1$, the inequality $v(B) \leq v(A)$ holds. Since both A and B are down-sets in (N, \preceq) , any such B is necessarily of the form $B = A \setminus \{i\}$, where i is a maximal element of A in (N, \preceq) . Note that $\downarrow i \subseteq A$ and $B \cap \downarrow i = (A \setminus \{i\}) \cap \downarrow i = \downarrow i$. Then supermodularity yields

$$v(A) = v(B \cup \downarrow i) \geq v(B) + v(\downarrow i) - v(\downarrow i).$$

Since v is 0-normalized and $\downarrow i \in \mathcal{J}$, Lemma 3.2 implies $v(\downarrow i) - v(\downarrow i) = 0$. \square

By the decomposition (3.2) we can now write $G_S(\mathcal{L})$ as the direct sum of cones,

$$G_S(\mathcal{L}) = G_S^*(\mathcal{L}) \oplus G_M(\mathcal{L}). \quad (3.3)$$

Specifically, the identity (3.3) means that $G_S(\mathcal{L}) = G_S^*(\mathcal{L}) + G_M(\mathcal{L})$ and $G_S^*(\mathcal{L}) \cap G_M(\mathcal{L}) = \{0\}$. Since $G_S^*(\mathcal{L})$ is a pointed polyhedral cone, it is generated by its finitely-many extreme rays.

In the next section we present a simple linear-algebraic criterion to test if a given 0-normalized supermodular game generates an extreme ray of $G_S^*(\mathcal{L})$. Our result automatically yields a criterion for extremality of games in $G_S(\mathcal{L})$: we say that a supermodular game $v \in G_S(\mathcal{L})$ is *extreme* if v^* generates an extreme ray of $G_S^*(\mathcal{L})$. Equivalently, $v \in G_S(\mathcal{L})$ is extreme if, and only if, the smallest face of $G_S(\mathcal{L})$ to which v belongs is an atom of the face lattice of $G_S(\mathcal{L})$. Indeed, faces of $G_S(\mathcal{L})$ are in one-to-one correspondence with faces of $G_S^*(\mathcal{L})$ by the relation $F = F^* + G_M(\mathcal{L})$, where F is a face of $G_S(\mathcal{L})$ and F^* a face of $G^*(\mathcal{L})$.

4. The cone of supermodular games

A *payoff vector* is any vector $x := (x_1, \dots, x_n) \in \mathbb{R}^n$. We define

$$x(A) := \sum_{i \in A} x_i, \quad \text{for any } A \in \mathcal{L},$$

and we always assume $x(\emptyset) := 0$. The *core* of $v \in G(\mathcal{L})$ is a convex polyhedron

$$C(v) := \{x \in \mathbb{R}^n \mid x(N) = v(N), x(A) \geq v(A) \text{ for each } A \in \mathcal{L}\}.$$

The elements of the core $\mathcal{C}(v)$ have the standard game-theoretic interpretation. Namely, no payoff vector $x \in \mathcal{C}(v)$ can be improved upon by any coalition $A \in \mathcal{L}$. In contrast with cores of games over Boolean lattices, the core of games over distributive lattices can be an unbounded polyhedron. In fact, assume $\mathcal{C}(v) \neq \emptyset$, where $v \in G(\mathcal{L})$. Then $\mathcal{C}(v)$ is bounded if and only if \mathcal{L} is a Boolean lattice; see [10, Chapter 3.3.3] or [5]. If $v \in G_S(\mathcal{L})$, then the polyhedron $\mathcal{C}(v)$ is pointed and its extreme points $\text{ext } \mathcal{C}(v)$ are characterized in Theorem 4.1 below.

Recall that we always assume that N is partially ordered by \preceq . In addition we also equip N with the total order of natural numbers \leq , so that (N, \leq) becomes a chain. We say that a permutation π of N is *compatible* with (N, \preceq) if π^{-1} is an order-preserving map from (N, \preceq) onto (N, \leq) . Here, the intended reading is that i is a rank of player $\pi(i)$. Define

$$\Pi_{\preceq} := \{\pi \mid \pi \text{ is a permutation compatible with } (N, \preceq)\}.$$

Compatible permutations are in bijection with maximal chains in \mathcal{L} . Put $A_0^\pi := \emptyset$ and $A_i^\pi := \{\pi(1), \dots, \pi(i)\}$ for each $i \in N$. Then, with each $\pi \in \Pi_{\preceq}$ we associate a maximal chain $\mathcal{C}^\pi := \{A_i^\pi \mid i \in N \cup \{0\}\}$. Conversely, starting from a maximal chain $\{A_0, \dots, A_n\}$ in \mathcal{L} , where $A_0 \subseteq \dots \subseteq A_n$, there is clearly a unique $\pi \in \Pi_{\preceq}$ such that $A_i = A_i^\pi$ for each $i \in N \cup \{0\}$.

A *marginal vector* of $v \in G(\mathcal{L})$ and $\pi \in \Pi_{\preceq}$ is the vector $x^{v, \pi} \in \mathbb{R}^n$ whose coordinates are defined as

$$x_{\pi(i)}^{v, \pi} := v(A_i^\pi) - v(A_{i-1}^\pi), \quad i \in N. \quad (4.1)$$

It follows directly from the definition of marginal vector that

$$v(A_i^\pi) = x^{v, \pi}(A_i^\pi), \quad \text{for all } \pi \in \Pi_{\preceq} \text{ and all } i \in N \cup \{0\}. \quad (4.2)$$

We will make an ample use of the following identity derived from (4.2):

$$v(A) = x^{v, \pi}(A), \quad \text{for all } \pi \in \Pi_{\preceq} \text{ and all } A \in \mathcal{C}^\pi. \quad (4.3)$$

For any $v \in G(\mathcal{L})$ and $\pi \in \Pi_{\preceq}$ we define

$$\mathcal{T}^\pi(v) := \{A \in \mathcal{L} \mid v(A) = x^{v, \pi}(A)\}.$$

Each coalition $A \in \mathcal{T}^\pi(v)$ is said to be *tight* with respect to v and π . Note that as a consequence of (4.3), the following inclusion holds:

$$\mathcal{C}^\pi \subseteq \mathcal{T}^\pi(v), \quad \text{for all } v \in G(\mathcal{L}) \text{ and all } \pi \in \Pi_{\preceq}. \quad (4.4)$$

Theorem 4.1. *Let \mathcal{L} be a finite distributive lattice and $v \in G(\mathcal{L})$. Then the following are equivalent:*

1. $v \in G_S(\mathcal{L})$.
2. $x^{v, \pi} \in \mathcal{C}(v)$, for each $\pi \in \Pi_{\preceq}$.
3. $\text{ext } \mathcal{C}(v) = \{x^{v, \pi} \mid \pi \in \Pi_{\preceq}\}$.
4. $v(A) = \min_{\pi \in \Pi_{\preceq}} x^{v, \pi}(A)$, for each $A \in \mathcal{L}$.

Proof. The equivalence of the first three items is well known; see [10, Theorem 3.27]. We show that 4. implies 2. Let $\sigma \in \Pi_{\preceq}$. Then, for all $A \in \mathcal{L}$,

$$x^{v,\sigma}(A) \geq \min_{\pi \in \Pi_{\preceq}} x^{v,\pi}(A) = v(A).$$

By (4.3) we have $x^{v,\sigma}(N) = v(N)$.

From 2. to 4. It suffices to show that for each $A \in \mathcal{L}$ there exists $\pi \in \Pi_{\preceq}$ such that $v(A) = x^{v,\pi}(A)$. Clearly, we can always find a maximal chain \mathcal{C}^π in \mathcal{L} such that $A \in \mathcal{C}^\pi$ for some $\pi \in \Pi_{\preceq}$. Then (4.3) yields $v(A) = x^{v,\pi}(A)$. \square

Remark 4.1. Many other characterizations of supermodularity can be found in the literature in case that \mathcal{L} is a Boolean lattice. See [21, Appendix A] for a comprehensive list of such conditions. In particular, the implication from 2. to 1. was proved by Ichiishi in [12]. The necessary and sufficient conditions involving specific marginal vectors can be found in [22].

Given $v \in G(\mathcal{L})$ let $\mathbf{x}^v: \Pi_{\preceq} \rightarrow \mathbb{R}^n$ be defined by

$$\mathbf{x}^v(\pi) := x^{v,\pi}, \quad \text{for all } \pi \in \Pi_{\preceq}.$$

Further, we consider a mapping $\mathbf{x}: G(\mathcal{L}) \rightarrow (\mathbb{R}^n)^{\Pi_{\preceq}}$ such that

$$\mathbf{x}(v) := \mathbf{x}^v, \quad \text{for all } v \in G(\mathcal{L}).$$

As in [21] we call \mathbf{x} the *payoff-array transformation*.

Lemma 4.1. *The payoff-array transformation \mathbf{x} is linear and injective.*

Proof. Linearity is a direct consequence of the identities $x^{v+w,\pi} = x^{v,\pi} + x^{w,\pi}$ and $x^{\alpha v,\pi} = \alpha x^{v,\pi}$, which are true for every $v, w \in G(\mathcal{L})$, all $\alpha \in \mathbb{R}$ and all $\pi \in \Pi_{\preceq}$. Assume that $v, w \in G(\mathcal{L})$ satisfy $\mathbf{x}^v = \mathbf{x}^w$ and let $A \in \mathcal{L}$. Then there exists a permutation $\pi \in \Pi_{\preceq}$ such that $A \in \mathcal{C}^\pi$. It follows from (4.3) and from the assumption that

$$v(A) = x^{v,\pi}(A) = x^{w,\pi}(A) = w(A).$$

Hence, \mathbf{x} is injective. \square

We describe the range of payoff-array transformation \mathbf{x} on the set of 0-normalized games. For any mapping $\mathbf{y}: \Pi_{\preceq} \rightarrow \mathbb{R}^n$ we denote $y^\pi := \mathbf{y}(\pi) \in \mathbb{R}^n$, for all $\pi \in \Pi_{\preceq}$.

Lemma 4.2. *Let $\mathbf{y}: \Pi_{\preceq} \rightarrow \mathbb{R}^n$. The following are equivalent:*

1. *There is a unique game $v \in G^*(\mathcal{L})$ such that $\mathbf{y} = \mathbf{x}^v$.*
2. *These conditions are satisfied:*

$$y^\pi(A) = y^\sigma(A) \quad \text{for all } \pi, \sigma \in \Pi_{\preceq} \text{ and all } A \in \mathcal{C}^\pi \cap \mathcal{C}^\sigma, \quad (\dagger)$$

$$y_i^\pi = 0 \quad \text{for all } \pi \in \Pi_{\preceq} \text{ and all } i \in N \text{ such that } \downarrow i \in \mathcal{C}^\pi. \quad (\dagger\dagger)$$

Proof. Let $\mathbf{y} = \mathbf{x}^v$ for some $v \in G^*(\mathcal{L})$. The equality in (\dagger) is a direct consequence of (4.3) since, for any $\pi, \sigma \in \Pi_{\leq}$ satisfying $A \in \mathcal{C}^\pi \cap \mathcal{C}^\sigma$, we get

$$y^\pi(A) = x^{v,\pi}(A) = v(A) = x^{v,\sigma}(A) = y^\sigma(A).$$

Further, let $\pi \in \Pi_{\leq}$ and $i \in N$ satisfy $\downarrow i \in \mathcal{C}^\pi$. Put $A := \downarrow i$ and observe that $A \in \mathcal{J}$ by Proposition 2.1. This implies that the unique predecessor of A in \mathcal{L} is $A^- = A \setminus \{i\}$ and $A^- \in \mathcal{C}^\pi$, by maximality of the chain \mathcal{C}^π . We obtain

$$y_i^\pi = x_i^{v,\pi} = x^{v,\pi}(A) - x^{v,\pi}(A^-) = v(A) - v(A^-) = 0,$$

where the third equality follows from (4.3) and the fourth one from 0-normalization of v (Lemma 3.2).

Conversely, assume that the conditions (\dagger) – $(\dagger\dagger)$ are true. If a game $v \in G^*(\mathcal{L})$ satisfying $\mathbf{y} = \mathbf{x}^v$ exists, then it is unique by injectivity of \mathbf{x} (Lemma 4.1). The condition (\dagger) guarantees that it is correct to define the game v as

$$v(A) := y^\pi(A) \quad \text{for all } \pi \in \Pi_{\leq} \text{ and all } A \in \mathcal{C}^\pi. \quad (4.5)$$

By the definition, $\mathbf{y} = \mathbf{x}^v$.

It remains to verify that v is 0-normalized. Let $A \in \mathcal{J}$. By Proposition 2.1 it follows that $A = \downarrow i$ for a unique $i \in N$. There exists some compatible permutation π satisfying $A \in \mathcal{C}^\pi$. Hence, by the definition of v and $(\dagger\dagger)$,

$$v(A) - v(A^-) = y^\pi(A) - y^\pi(A^-) = y_i^\pi = 0.$$

This means that v is 0-normalized and the proof is finished. \square

Remark 4.2. A mapping $\mathbf{y}: \Pi_{\leq} \rightarrow \mathbb{R}^n$, whose special case is the payoff-array transformation \mathbf{x} , can be viewed as a finite collection of possibly repeating points in \mathbb{R}^n labeled by permutations. This interpretation appears in [3], where a map \mathbf{y} from a finite set into \mathbb{R}^n is termed a *point configuration*.

5. Main result

Denote

$$N^\pi(v) := \{i \in N \mid x_i^{v,\pi} = 0\}.$$

The main theorem gives a simple criterion how to recognize extreme games among all 0-normalized supermodular games.

Theorem 5.1. *Let $v \in G_S^*(\mathcal{L})$ be nonzero. Then the following are equivalent:*

1. *The game v is extreme in $G_S^*(\mathcal{L})$.*
2. *If $\mathbf{y}: \Pi_{\leq} \rightarrow \mathbb{R}^n$ satisfies the conditions*

$$y^\pi(A) = y^\sigma(A) \quad \text{for all } \pi, \sigma \in \Pi_{\leq} \text{ and all } A \in \mathcal{T}^\pi(v) \cap \mathcal{T}^\sigma(v), \quad (*)$$

$$y_i^\pi = 0 \quad \text{for all } \pi \in \Pi_{\leq} \text{ and all } i \in N^\pi(v), \quad (**)$$

then $\mathbf{y} = \alpha \mathbf{x}^v$, for some $\alpha \in \mathbb{R}$.

We will briefly comment on the interpretation of Theorem 5.1. The linear-algebraic characterization of extreme rays (Minkowski-Weyl-Farkas theorem) provides a universally valid method for any pointed polyhedral cone. Our approach based on Theorem 5.1 uses the structure of the objects we are dealing with (supermodular functions on finite distributive lattices). Hence, one need not to assemble all the relevant linear systems “blindly” using the linear-algebraic technique. Indeed, the shape of such systems is fully determined by the combinatorial core structure, which is the collection of tight sets and null sets. Loosely speaking, a supermodular game is extreme if, and only if, the core structure is rigid in sense that the associated linear system is uniquely solvable.

We prepare a lemma to be used in the proof of Theorem 5.1. For any $v \in G(\mathcal{L})$, put

$$\mathcal{F}_v := \{\{A, B\} \subseteq \mathcal{L} \mid v(A \cup B) + v(A \cap B) = v(A) + v(B), A \parallel B\}. \quad (5.1)$$

where

$$A \parallel B \text{ means } A \not\subseteq B \text{ and } B \not\subseteq A.$$

For any point configuration $\mathbf{y} : \Pi_{\leq} \rightarrow \mathbb{R}^n$ and a game $v \in G(\mathcal{L})$, we consider the following property:

$$\begin{aligned} y^\pi(A) = y^\sigma(A), \quad \text{for each } \{A, B\} \in \mathcal{F}_v \text{ and all } \pi, \sigma \in \Pi_{\leq} \\ \text{such that } A \cap B, B, A \cup B \in \mathcal{C}^\pi \text{ and } A \in \mathcal{C}^\sigma. \end{aligned} \quad (5.2)$$

Lemma 5.1. *Let $v \in G_S^*(\mathcal{L})$ and let $\mathbf{y} : \Pi_{\leq} \rightarrow \mathbb{R}^n$ be such that (*) and (**) are satisfied. Then \mathbf{y} fulfills †, ††, and (5.2).*

Proof. Assume that \mathbf{y} satisfies (*) and (**). It is easy to see that † and †† are true. In order to prove (5.2), let $\{A, B\} \in \mathcal{F}_v$, $\pi, \sigma \in \Pi_{\leq}$, and $A \cap B, B, A \cup B \in \mathcal{C}^\pi$, $A \in \mathcal{C}^\sigma$. Since

$$v(A) = v(A \cup B) + v(A \cap B) - v(B) = x^{v, \pi}(A \cup B) + x^{v, \pi}(A \cap B) - x^{v, \pi}(B) = x^{v, \pi}(A),$$

we get $A \in \mathcal{T}^\pi(v)$. Hence, $A \in \mathcal{T}^\pi(v) \cap \mathcal{C}^\sigma$ and (*) says that $y^\pi(A) = y^\sigma(A)$. \square

Proof. (of Theorem 5.1) Let $v \in G_S^*(\mathcal{L})$ be nonzero. We need to show that v is extreme if and only if the following inclusion holds true:

$$\{\mathbf{y} : \Pi_{\leq} \rightarrow \mathbb{R}^n \mid \mathbf{y} \text{ satisfies } (*), (**)\} \subseteq \{\alpha x^v \mid \alpha \in \mathbb{R}\}. \quad (5.3)$$

By the Minkowski-Weyl-Farkas theorem (see [1, Theorem 3.34]), v is extreme if and only if v belongs to the one-dimensional solution space of some set of tight inequalities for $G_S^*(\mathcal{L})$ of the form $w(A \cup B) + w(A \cap B) - w(A) - w(B) \geq 0$, for all $A, B \in \mathcal{L} \setminus \{\emptyset, N\}$. Define

$$G(v) := \{w \in G^*(\mathcal{L}) \mid w(A \cup B) + w(A \cap B) = w(A) + w(B), \text{ for all } \{A, B\} \in \mathcal{F}_v\},$$

where \mathcal{F}_v is as in (5.1). Thus, extremality of v is equivalent to the condition

$$G(v) = \{\alpha v \mid \alpha \in \mathbb{R}\}. \quad (5.4)$$

Putting $\mathbf{x}(G(v)) := \{\mathbf{x}^w \mid w \in G(v)\}$ and using Lemma 4.1, it is immediate that (5.4) holds if and only if

$$\mathbf{x}(G(v)) = \{\alpha x^v \mid \alpha \in \mathbb{R}\}. \quad (5.5)$$

We claim that

$$\mathbf{x}(G(v)) \supseteq \{\mathbf{y} : \Pi \rightarrow \mathbb{R}^n \mid \mathbf{y} \text{ satisfies } (\dagger), (\dagger\dagger), \text{ and } (5.2)\}. \quad (5.6)$$

Let \mathbf{y} satisfies (\dagger) , $(\dagger\dagger)$, and (5.2). Lemma 4.2 provides a unique $w \in G^*(\mathcal{L})$ such that $\mathbf{y} = \mathbf{x}^w$. We need to verify that $w \in G(v)$. To this end, let $\{A, B\} \in \mathcal{F}_v$. Pick permutations $\pi, \sigma \in \Pi_{\leq}$ such that $A \cap B, B, A \cup B \in \mathcal{C}^\pi$ and $A \in \mathcal{C}^\sigma$. Then (5.2) shows that

$$w(A \cup B) + w(A \cap B) - w(B) = y^\pi(A) = y^\sigma(A) = w(A).$$

Hence, $w \in G(v)$. Finally, from (5.5), (5.6), and Lemma 5.1 we get (5.3), and the proof is finished. \square

We will apply Theorem 5.1 to the cone of supermodular games on the distributive lattice \mathcal{L} from Example 2.1. The computations were carried out in the package Convex for Maple [7].

Example 5.1. The cone $G_S^*(\mathcal{L})$ is embedded into \mathbb{R}^9 and its dimension is 5. It has 6 extreme rays. We will enumerate their minimal integer generators. The parentheses and commas are omitted for the sake of brevity in what follows. Whenever $v_i(A)$ is missing, we put $v_i(A) := 0$.

- $v_1(24) = v_1(234) = v_1(N) = 1$.
- $v_2(34) = v_2(234) = v_2(N) = 1$.
- $v_3(23) = v_3(123) = v_3(234) = v_3(N) = 1$.
- $v_4(234) = v_4(N) = 1$.
- $v_5(23) = v_5(24) = v_5(34) = v_5(123) = 1$, $v_5(234) = v_5(N) = 2$.
- $v_6(N) = 1$.

We will check that v_1 is extreme using Theorem 5.1. Since there are 8 maximal chains in \mathcal{L} , there are 8 compatible permutations: $\pi_1 = (2314)$, $\pi_2 = (2341)$, $\pi_3 = (2431)$, $\pi_4 = (3241)$, $\pi_5 = (3241)$, $\pi_6 = (3421)$, $\pi_7 = (4231)$, and $\pi_8 = (4321)$. Let $I_1 := \{1, \dots, 5\}$ and $I_2 := \{6, 7, 8\}$. There are only 2 marginal vectors associated with v_1 ,

$$x^{v_1, \pi_i} = \begin{cases} (0, 0, 0, 1) & i \in I_1, \\ (0, 1, 0, 0) & i \in I_2. \end{cases}$$

This means that the tight sets are

$$\mathcal{T}^{\pi_i}(v_1) = \begin{cases} \{\emptyset, 2, 3, 23, 24, 123, 234, N\} & i \in I_1, \\ \{\emptyset, 3, 4, 24, 34, 234, N\} & i \in I_2. \end{cases}$$

Hence, the conditions (*) and (**) for $\mathbf{y}: \Pi_{\leq} \rightarrow \mathbb{R}^4$ are in the form of linear equalities, for all $i \in I_1$ and $j \in I_2$:

$$y_1^{\pi_i} = y_2^{\pi_i} = y_3^{\pi_i} = 0 \quad (5.7)$$

$$y_1^{\pi_j} = y_3^{\pi_j} = y_4^{\pi_j} = 0 \quad (5.8)$$

$$y_3^{\pi_i} = y_3^{\pi_j} \quad (5.9)$$

$$y_2^{\pi_i} + y_4^{\pi_i} = y_2^{\pi_j} + y_4^{\pi_j} \quad (5.10)$$

$$y_2^{\pi_i} + y_3^{\pi_i} + y_4^{\pi_i} = y_2^{\pi_j} + y_3^{\pi_j} + y_4^{\pi_j} \quad (5.11)$$

$$y_1^{\pi_i} + y_2^{\pi_i} + y_3^{\pi_i} + y_4^{\pi_i} = y_1^{\pi_j} + y_2^{\pi_j} + y_3^{\pi_j} + y_4^{\pi_j} \quad (5.12)$$

The linear system above has a unique solution up to a real multiple. Observe that $y_4^{\pi_i} = y_2^{\pi_j}$, for all $i \in I_1$ and $j \in I_2$, as a consequence of (5.7), (5.8), and (5.10). Let $\alpha \in \mathbb{R}$. Then necessarily $\mathbf{y} = \alpha \mathbf{x}^{v_1}$. Thus, v_1 is extreme by Theorem 5.1.

Remark 5.1. It is natural to ask for a game-theoretic meaning of the extreme supermodular games. Since the supermodular cone is finitely-generated, every supermodular game is a conic combination of the extreme ones. There are important solution concepts [14], such as the core or Shapley value, which are linear maps on the supermodular cone. Hence, such solution concepts preserve every conic combination of supermodular games. From this viewpoint, extreme supermodular games play the role of basic building block since they fully determine values of any linear solution concept on the supermodular cone.

6. Faces and core structure

Let $\Phi(G_S(\mathcal{L}))$ be the *face lattice* of $G_S(\mathcal{L})$, that is, the family of all nonempty faces of $G_S(\mathcal{L})$ ordered by inclusion \subseteq . In what follows we will describe the structure of this face lattice. For any subset $G \subseteq G_S(\mathcal{L})$ we define

$$[G] := \bigcap \{F \in \Phi(G_S(\mathcal{L})) \mid F \supseteq G\},$$

the smallest face containing G . Join \vee and meet \wedge in $\Phi(G_S(\mathcal{L}))$ are computed as

$$F \sqcup G = [F \cup G] \text{ and } F \sqcap G = F \cap G, \quad \text{for all } F, G \in \Phi(G_S(\mathcal{L})).$$

For any face F , let $\text{relint } F$ be the relative interior of F . Put $\Phi'(G_S(\mathcal{L})) = \{\text{relint } F \mid F \in \Phi(G_S(\mathcal{L}))\}$. Then $\Phi'(G_S(\mathcal{L}))$ is a lattice isomorphic to $\Phi(G_S(\mathcal{L}))$ in which the top is $\text{relint } G_S(\mathcal{L})$, the bottom is \emptyset , and the join and the meet are given by

$$\begin{aligned} \text{relint } F \vee \text{relint } G &= \text{relint}(F \sqcup G), \\ \text{relint } F \wedge \text{relint } G &= \text{relint}(F \sqcap G). \end{aligned}$$

The following lemma describes the relation between tight sets of $v \in G_S(\mathcal{L})$ and faces of $G_S(\mathcal{L})$.

Lemma 6.1. *Let $v \in G_S(\mathcal{L})$. The following holds.*

1. Let $\pi \in \Pi_{\leq}$ and $A, B \in \mathcal{T}^{\pi}(v)$, $A \parallel B$. Then $\{A, B\} \in \mathcal{F}_v$.

2. Let $\{A, B\} \in \mathcal{F}_v$ and $\pi \in \Pi_{\leq}$ such that $B, A \cup B, A \cap B \in \mathcal{C}^\pi$. Then $A \in \mathcal{T}^\pi(v) \setminus \mathcal{C}^\pi$.

Proof. 1. Let $v \in G_S(\mathcal{L})$, $\pi \in \Pi_{\leq}$, and $A, B \in \mathcal{T}^\pi(v)$ with $A \parallel B$. It is well known that $\mathcal{T}^\pi(v)$ is closed under union and intersection for supermodular games. Hence, the equality $x^{v,\pi}(A) + x^{v,\pi}(B) = x^{v,\pi}(A \cup B) + x^{v,\pi}(A \cap B)$ yields $v(A) + v(B) = v(A \cup B) + v(A \cap B)$. Therefore, $\{A, B\} \in \mathcal{F}_v$.

2. Let $\{A, B\} \in \mathcal{F}_v$ and let $\pi \in \Pi_{\leq}$ be such that $B, A \cup B, A \cap B \in \mathcal{C}^\pi$. Then

$$v(A) = v(A \cup B) + v(A \cap B) - v(B) = x^{v,\pi}(A \cup B) + x^{v,\pi}(A \cap B) - x^{v,\pi}(B) = x^{v,\pi}(A).$$

Since $A \parallel B$, we get $A \in \mathcal{T}^\pi(v) \setminus \mathcal{C}^\pi$. \square

Games in $G_S(\mathcal{L})$ belong to the same face if and only if they possess identical structure of their tight sets. Precisely:

Proposition 6.1. *Let $v, w \in G_S(\mathcal{L})$. The following are equivalent.*

1. $v, w \in \text{relint } F$, for some $F \in \Phi(G_S(\mathcal{L}))$.
2. $\mathcal{T}^\pi(v) = \mathcal{T}^\pi(w)$, for every $\pi \in \Pi_{\leq}$.

Proof. Let $v, w \in \text{relint } F$ for some $F \in \Phi(G_S(\mathcal{L}))$. If they are linearly dependent, then the statement is trivial. Assume that v and w are linearly independent and let L be the unique line in the linear space $G(\mathcal{L})$ such that $v, w \in L$. Then $L \cap (F \setminus \text{relint } F) = \{u, u'\}$, where $u \neq u'$. Since $v, w \in \text{relint } F$ there exist $0 < \alpha, \beta < 1$ such that $v = \alpha u + (1 - \alpha)u'$ and $w = \beta u + (1 - \beta)u'$.

Let $\pi \in \Pi_{\leq}$ and $A \in \mathcal{T}^\pi(v)$. By linearity (Lemma 4.1) we get

$$\alpha x^{u,\pi}(A) + (1 - \alpha)x^{u',\pi}(A) = x^{v,\pi}(A) = v(A) = \alpha u(A) + (1 - \alpha)u'(A).$$

Since $x^{u,\pi} \in \mathcal{C}(u)$ and $x^{u',\pi} \in \mathcal{C}(u')$, this implies $x^{u,\pi}(A) = u(A)$ and $x^{u',\pi}(A) = u'(A)$, which means that $A \in \mathcal{T}^\pi(u) \cap \mathcal{T}^\pi(u')$. Hence,

$$w(A) = \beta u(A) + (1 - \beta)u'(A) = \beta x^{u,\pi}(A) + (1 - \beta)x^{u',\pi}(A) = x^{w,\pi}(A).$$

This proves the inclusion $\mathcal{T}^\pi(v) \subseteq \mathcal{T}^\pi(w)$. The opposite inclusion is established analogously.

To prove the converse, assume $\mathcal{T}^\pi(v) = \mathcal{T}^\pi(w)$ for all $\pi \in \Pi_{\leq}$. It suffices to show that $\mathcal{F}_v = \mathcal{F}_w$, where \mathcal{F}_v is as in (5.1), since this already implies existence of a unique $F \in \Phi(G_S(\mathcal{L}))$ such that $v, w \in \text{relint } F$. First, we prove

$$\mathcal{F}_v \subseteq \mathcal{F}_w. \tag{6.1}$$

Let $\{A, B\} \in \mathcal{F}_v$. There exists $\pi \in \Pi_{\leq}$ such that $A \cap B, A, A \cup B \in \mathcal{C}^\pi$. Hence, $A \in \mathcal{T}^\pi(v)$ and

$$\begin{aligned} v(B) &= v(A \cup B) + v(A \cap B) - v(A) \\ &= x^{v,\pi}(A \cup B) + x^{v,\pi}(A \cap B) - x^{v,\pi}(A) = x^{v,\pi}(B), \end{aligned}$$

which means $B \in \mathcal{T}^\pi(v) = \mathcal{T}^\pi(w)$. Then $A, B \in \mathcal{T}^\pi(w)$, and by Lemma 6.1 (1) $\{A, B\} \in \mathcal{F}_w$, so (6.1) holds. The proof of inclusion $\mathcal{F}_w \subseteq \mathcal{F}_v$ is analogous. \square

Let $\mathcal{S}(\mathcal{L})$ be the lattice of all sublattices of \mathcal{L} ordered by set inclusion \subseteq . The *core structure* of $v \in G_S(\mathcal{L})$ (cf. [13] and [21, Definition 4]) is the mapping $\mathcal{T}(v): \Pi_{\leq} \rightarrow \mathcal{S}(\mathcal{L})$ defined as

$$\pi \in \Pi_{\leq} \mapsto \mathcal{T}^\pi(v).$$

The above definition is correct since $\mathcal{T}^\pi(v)$ is a lattice as a consequence of Lemma 6.1. By Proposition 6.1, $\mathcal{T}(v) = \mathcal{T}(w)$ for all $v, w \in \text{relint } F$. Hence, we may define a mapping

$$\mathcal{T}: \Phi(G_S(\mathcal{L})) \rightarrow \mathcal{S}(\mathcal{L})^{\Pi_{\leq}}$$

by

$$\mathcal{T}(F) := \mathcal{T}(v), \quad \text{for any } v \in \text{relint } F.$$

We will order the elements of $\mathcal{S}(\mathcal{L})^{\Pi_{\leq}}$ by the product order \subseteq inherited from $\mathcal{S}(\mathcal{L})$. Specifically, for any $\mathcal{U}, \mathcal{V} \in \mathcal{S}(\mathcal{L})^{\Pi_{\leq}}$,

$$\mathcal{U} \subseteq \mathcal{V} \text{ whenever } \mathcal{U}^\pi \subseteq \mathcal{V}^\pi, \quad \text{for all } \pi \in \Pi_{\leq}.$$

Proposition 6.2. *The mapping \mathcal{T} is injective, order-reversing, and its inverse \mathcal{T}^{-1} is also order-reversing.*

Proof. \mathcal{T} injective is an easy consequence of Proposition 6.1. We will prove that \mathcal{T} is an order-reversing map. Let $F_1 \subseteq F_2$ be faces of $G_S(\mathcal{L})$ and select arbitrarily $v_1 \in \text{relint } F_1$ and $v_2 \in \text{relint } F_2$. We want to show

$$\mathcal{T}^\pi(v_2) \subseteq \mathcal{T}^\pi(v_1) \quad \text{for every } \pi \in \Pi_{\leq}. \quad (6.2)$$

Let $\pi \in \Pi_{\leq}$ and $A \in \mathcal{T}^\pi(v_2)$. Using (4.1), this is equivalent to saying that v_2 is a solution of the equation in v :

$$\sum_{i \in A} (v(A_{\pi^{-1}(i)}^\pi) - v(A_{\pi^{-1}(i-1)}^\pi)) = v(A). \quad (6.3)$$

As this equation is satisfied by all games in $\text{relint } F_2$ and only these ones, it follows that (6.3) is implied by the equalities determining $\text{relint } F_2$, that is, those corresponding to \mathcal{F}_{v_2} . As $F_1 \subseteq F_2$, $\text{relint } F_1$ is determined by a superset of equalities, and therefore the equality (6.3) is also satisfied by v_1 . Hence $A \in \mathcal{T}^\pi(v_1)$.

To show that \mathcal{T}^{-1} is order-reversing, let $\mathcal{T}^\pi(v_1) \subseteq \mathcal{T}^\pi(v_2)$ for all $\pi \in \Pi_{\leq}$, where $v_1 \in \text{relint } F_1$ and $v_2 \in \text{relint } F_2$, for some faces F_1 and F_2 . We will prove that $F_2 \subseteq F_1$, which is the same as $\mathcal{F}_{v_1} \subseteq \mathcal{F}_{v_2}$. Let $\{A, B\} \in \mathcal{F}_{v_1}$. Then there exists π s.t. $A \cap B, A, A \cup B \in \mathcal{C}^\pi$. Hence, by Lemma 6.1 (2), $B \in \mathcal{T}^\pi(v_1) \setminus \mathcal{C}^\pi$, hence $B \in \mathcal{T}^\pi(v_2) \setminus \mathcal{C}^\pi$. As $A \in \mathcal{C}^\pi \subseteq \mathcal{T}^\pi(v_2)$ and $A \parallel B$, by Lemma 6.1 (1), $\{A, B\} \in \mathcal{F}_{v_2}$. \square

Corollary 6.1. *\mathcal{T} is a lattice isomorphism from the face lattice $\Phi(G_S(\mathcal{L}))$ onto a sublattice of $\mathcal{S}(\mathcal{L})^{\Pi_{\leq}}$.*

Remark 6.1. The same reasoning can be applied to the face lattice of all 0-normalized supermodular games, $\Phi(G_S^*(\mathcal{L}))$. Indeed, it follows from the direct sum decomposition (3.3) that $\Phi(G_S^*(\mathcal{L}))$ and $\Phi(G_S(\mathcal{L}))$ are isomorphic lattices.

7. Facets of the cone of supermodular games

Kuipers et al. [13, Corollary 11] characterised those supermodular inequalities which determine the facets of $G_S(\mathcal{L})$ when $\mathcal{L} = 2^N$. Specifically, the facet-determining inequalities are

$$v(A \cup \{i, j\}) - v(A \cup \{i\}) - v(A \cup \{j\}) + v(A) \geq 0 \quad (7.1)$$

for all $A \subseteq N$ and every pair of distinct $i, j \in N \setminus A$. The next theorem identifies the facets of the cone $G_S(\mathcal{L})$, for any \mathcal{L} . Since \mathcal{L} is the lattice of down-sets of a poset (N, \preceq) , if $A \in \mathcal{L}$ and $i \in N$ are such that $\downarrow i \subseteq A$, then $A \cup \{i\} \in \mathcal{L}$.

Theorem 7.1. *The facets of $G_S(\mathcal{L})$ are given by the inequalities of the form*

$$v(A \cup \{i, j\}) - v(A \cup \{i\}) - v(A \cup \{j\}) + v(A) \geq 0 \quad (7.2)$$

with $A \in \mathcal{L}$ and distinct $i, j \in N \setminus A$ such that $\downarrow i \subseteq A$ and $\downarrow j \subseteq A$.

Proof. Observe that, since the lattice \mathcal{L} is ranked¹, if A and $A \cup B$ with $B \subseteq N \setminus A$ are in \mathcal{L} , then $A \cup \{i\} \in \mathcal{L}$ for some $i \in B$.

First, we show that any supermodular inequality can be derived from those of type (7.2). Consider

$$v(A \cup B) - v(A) - v(B) + v(A \cap B) \geq 0, \quad (A, B \in \mathcal{L}) \quad (7.3)$$

with $|A \Delta B| > 2$. We show by induction on $|A \Delta B|$ that (7.3) can be derived from (7.2). First we establish the result for $|A \Delta B| = 3$. Take

$$v(A \cup \{i, j, k\}) - v(A \cup \{i, j\}) - v(A \cup \{k\}) + v(A) \geq 0, \quad (7.4)$$

with $A \cup \{i, j\}, A \cup \{k\} \in \mathcal{L}$. Supposing $A' := A \cup \{i\} \in \mathcal{L}$ (by the preliminary remark, either $A \cup \{i\}$ or $A \cup \{j\}$ belongs to \mathcal{L}), we have by the assumption

$$\begin{aligned} v(A' \cup \{j, k\}) - v(A' \cup \{j\}) - v(A' \cup \{k\}) + v(A') &\geq 0, \\ v(A \cup \{i, k\}) - v(A \cup \{i\}) - v(A \cup \{k\}) + v(A) &\geq 0, \end{aligned}$$

whose sum yields (7.4).

We suppose that the result holds for $|A \Delta B| = k$, where $k \geq 3$, and let us prove it for $|A \Delta B| = k + 1$. Consider the inequality

$$v(A \cup \{i_1, \dots, i_\ell, \dots, i_{k+1}\}) - v(A \cup \{i_1, \dots, i_\ell\}) - v(A \cup \{i_{\ell+1}, \dots, i_{k+1}\}) + v(A) \geq 0, \quad (7.5)$$

with $A \cup \{i_1, \dots, i_\ell\}, A \cup \{i_{\ell+1}, \dots, i_{k+1}\} \in \mathcal{L}$. Put $A' := A \cup \{i_{\ell+1}\} \in \mathcal{L}$. It follows from the assumption that

$$\begin{aligned} v(A' \cup \{i_1, \dots, i_\ell, i_{\ell+2}, \dots, i_{k+1}\}) - v(A' \cup \{i_1, \dots, i_\ell\}) - v(A' \cup \{i_{\ell+2}, \dots, i_{k+1}\}) + v(A') &\geq 0 \\ v(A \cup \{i_1, \dots, i_{\ell+1}\}) - v(A \cup \{i_{\ell+1}\}) - v(A \cup \{i_1, \dots, i_\ell\}) + v(A) &\geq 0, \end{aligned}$$

whose sum gives (7.5).

¹A *ranked* (also *graded*) *lattice* is a lattice in which all the maximal chains have the same cardinality.

Second, we prove that no inequality of type (7.2) is redundant. It is clearly sufficient to prove the result for the Boolean lattice $\mathcal{L} = 2^N$. Consider (7.2) for fixed A, i, j and define the following game $v_{A,i,j}$ for an arbitrary $\varepsilon > 0$:

$$v_{A,i,j}(B) := \begin{cases} \varepsilon & \text{if } B = A \cup \{i\}, \\ 0 & \text{if } B \subset A \cup \{i\} \text{ or } B = A \cup \{j\} \text{ or } B \supset A \cup \{i\}, \\ -\varepsilon & \text{otherwise.} \end{cases}$$

We claim that $v_{A,i,j}$ satisfies all the inequalities (7.2) except the one with A, i, j . This proves that the latter inequality is not redundant.

Proof of the claim. Put $v := v_{A,i,j}$, $i := 1, j := 2$. Clearly,

$$v(A \cup \{1, 2\}) - v(A \cup \{1\}) - v(A \cup \{2\}) + v(A) = -\varepsilon < 0.$$

Consider the quantity

$$\Delta(B, i, j) := v(B \cup \{i, j\}) - v(B \cup \{i\}) - v(B \cup \{j\}) + v(B).$$

We first study $\Delta(B, i, j)$ when one of the terms has value ε . If $v(B \cup \{i, j\}) = \varepsilon$ or $v(B) = \varepsilon$, the other terms can be 0 or $-\varepsilon$, so that $\Delta(B, i, j) \geq 0$. If $v(B \cup \{i\}) = \varepsilon$ (i.e., $B = A, i = 1$), then $v(B \cup \{j\}) = -\varepsilon$, unless $j = 2$. Supposing this is not the case, we have then $v(B \cup \{i, j\}) = v(B) = 0$. Hence, $\Delta(B, i, j) = 0$.

We study now $\Delta(B, i, j)$ when the values of the terms are either 0 or $-\varepsilon$. Suppose that $v(B \cup \{i\}) = v(B \cup \{j\}) = 0$. Both $B \cup \{i\}$ and $B \cup \{j\}$ are either proper subsets of $A \cup \{1\}$ or proper supersets of it. Then both B and $B \cup \{i, j\}$ are either subsets or supersets of $A \cup \{1\}$, and the equality is impossible since we excluded the value ε for v . Therefore, $v(B \cup \{i, j\}) = v(B) = 0$, and $\Delta(B, i, j) = 0$. Suppose now that $v(B \cup \{i\}) = 0$ and $v(B \cup \{j\}) = -\varepsilon$. Then either $B \cup \{i\} \subset A \cup \{1\}$ or $B \cup \{i\} \supset A \cup \{1\}$, and in both cases $j \notin A \cup \{1\}$. It follows that, if $B \cup \{i\} \subset A \cup \{1\}$, then $B \subset A \cup \{1\}$, therefore $v(B) = 0$, yielding $\Delta(B, i, j) \geq 0$. If $B \cup \{i\} \supset A \cup \{1\}$, then $v(B \cup \{i, j\}) = 0$, hence the same conclusion holds. Finally, suppose $v(B \cup \{i\}) = v(B \cup \{j\}) = -\varepsilon$. Then, in any case, $\Delta(B, i, j) \geq 0$. \square

Example 7.1. Let (N, \preceq) and \mathcal{L} be as in Example 2.1. Applying Theorem 7.1 we see that the facets of $G_S(\mathcal{L})$ are in bijection with the linear inequalities of the form

1. $v(\{i, j\}) \geq v(\{i\}) + v(\{j\})$, for all distinct $i, j \in N \setminus \{1\}$,
2. $v(A \cup B) + v(A \cap B) \geq v(A) + v(B)$, for all distinct $A, B \in \mathcal{L}$ satisfying $|A| = |B| = 2$,
3. $v(N) + v(\{2, 3\}) \geq v(\{1, 2, 3\}) + v(\{2, 3, 4\})$.

Thus, there are 7 facets, whereas the cone $G_S(2^N)$ of supermodular games on the Boolean lattice 2^N has $\binom{n}{2} \cdot 2^{n-2} = 24$ facets by (7.1).

Acknowledgements

The work of Tomáš Kroupa has been supported from the Czech Science Foundation project GA16-12010S.

[1] C. Aliprantis, R. Tourky, *Cones and duality*, volume 84 of *Graduate Studies in Mathematics*, American Mathematical Society, Providence, RI, 2007.

- [2] G. Birkhoff, *Lattice Theory*, American Mathematical Society, 1948.
- [3] J. De Loera, J. Rambau, F. Santos, *Triangulations. Structures for Algorithms and Applications*, volume 25 of *Algorithms and Computation in Mathematics*, Springer-Verlag Berlin Heidelberg, 2010.
- [4] R. van den Brink, R. Gilles, Axiomatizations of the conjunctive permission value for games with permission structures, *Games and Economic Behavior*, 12:113–126, 1996.
- [5] J. J. Derks, R. P. Giles, Hierarchical organization structures and constraints on coalition formation, *International Journal of Game Theory*, 24(2), 147–163.
- [6] U. Faigle, W. Kern, The Shapley value for cooperative games under precedence constraints, *International Journal of Game Theory*, 21:249–266, 1992.
- [7] M. Franz, Convex – a Maple package for convex geometry, version 1.2.0, Available at <http://www.math.uwo.ca/faculty/franz/convex/>, 2016.
- [8] S. Fujishige, *Submodular functions and optimization*, volume 58 of *Annals of Discrete Mathematics*, Elsevier B. V., Amsterdam, second edition, 2005.
- [9] M. Grabisch, The core of games on ordered structures and graphs, *Annals of Operations Research*, 204:33–64, 2013.
- [10] M. Grabisch, *Set functions, games and capacities in decision making*, Springer, 2016.
- [11] M. Grabisch, L. J. Xie, The restricted core of games on distributive lattices: how to share benefits in a hierarchy, *Mathematical Methods of Operations Research*, 73:189–208, 2011.
- [12] T. Ichiishi, Super-modularity: Applications to convex games and to the greedy algorithm for LP, *Journal of Economic Theory*, 25(2):283–286, 1981.
- [13] J. Kuipers, D. Vermeulen, M. Voorneveld, A generalization of the Shapley–Ichiishi result, *International Journal of Game Theory*, 39(4):585–602, 2010.
- [14] B. Peleg, P. Sudhölter, *Introduction to the theory of cooperative games*, volume 34 of *Theory and Decision Library. Series C: Game Theory, Mathematical Programming and Operations Research*, Springer, Berlin, second edition, 2007.
- [15] G.-C. Rota, On the foundations of combinatorial theory. I. Theory of Möbius functions, *Zeitschrift für Wahrscheinlichkeitstheorie und Verwandte Gebiete*, 2:340–368 (1964), 1964.
- [16] G.-C. Rota, On the combinatorics of the Euler characteristic, In *Studies in Pure Mathematics (Presented to Richard Rado)*, pages 221–233. Academic Press London, 1971.
- [17] J. Rosenmüller, H. G. Weidner, Extreme convex set functions with finite carrier: general theory, *Discrete Mathematics*, 10:343–382, 1974.
- [18] D. Schmeidler, Subjective Probability and Expected Utility without Additivity, *Econometrica*, 57:571–587, 1989.
- [19] L. S. Shapley, Cores of convex games, *International Journal of Game Theory*, 1:11–26, 1971.
- [20] R. P. Stanley, *Enumerative Combinatorics, Vol. 1*, Cambridge University Press, second edition, 2012.
- [21] M. Studený, T. Kroupa, Core-based criterion for extreme supermodular games, *Discrete Applied Mathematics*, 206:122–151, 2016.
- [22] B. van Velzen, H. Hamers, H. Norde, Characterizing convexity of games using marginal vectors, *Discrete Applied Mathematics*, 143:298–306, 2004.