

Worldwide estimation of parameters for a simple reaction-diffusion model of urban growth

J. Raimbault^{1,2,3*}


*j.raimbault@ucl.ac.uk

¹CASA, UCL

²UPS CNRS 3611 Complex Systems Institute Paris

³UMR CNRS 8504 Géographie-cités

International Land-Use Symposium 2019
Session Drivers, Mechanisms, Tools
5th December 2019





- Cellular automata models or models of land-use change generally highly data-driven
- May be case-specific or conclusions difficult to generalize (see Sleuth for example [Clarke et al., 2007])
- Issues to calibrate and validate these models [Engelen and White, 2008]

Urban morphogenesis models approach:

- *At the crossroad between Urban Simulation and Artificial Life, stylized models of urban growth processes*
- *Importance of parsimony: generative modeling as a tool to understand processes*

→ Explore a simple model to capture morphogenesis based on abstract representation of urban processes; test their ability to reproduce existing urban systems.

→ The simple reaction-diffusion model of urban growth is calibrated worldwide using GHSL, and in time.

→ Globally comparable insights into processes of concentration and dispersion.

- Crucial role of the interplay between concentration forces and dispersion forces [Fujita and Thisse, 1996] in keeping Urban Systems at the border of chaos
- Potentiality of aggregation mechanisms (such as Simon model) to produce power laws [Dodds et al., 2017]
- Link with Reaction-diffusion approaches in Morphogenesis [Turing, 1952]
- Extension of a DLA-type model introduced by [Batty, 1991], with simple abstract processes of population aggregation and diffusion

Raimbault, J. (2018). Calibration of a density-based model of urban morphogenesis. PloS one, 13(9), e0203516.

→ Grid world with cell populations $(P_i(t))_{1 \leq i \leq N^2}$.

→ At each time step:

- 1** Population growth with exogenous rate N_G , attributed independently to a cell following a preferential attachment of strength α
- 2** Population is diffused n_d times with strength β

→ Stopping criterion: fixed maximal population P_m .

→ Output measured by morphological indicators: Moran index, average distance, rank-size hierarchy, entropy.

- 1 Rank-size slope γ , given by $\ln(P_i/P_0) \sim k + \gamma \cdot \ln(\tilde{i}/i_0)$ where \tilde{i} are the indexes of the distribution sorted in decreasing order.
- 2 Entropy of the distribution:

$$\mathcal{E} = \sum_{i=1}^M \frac{P_i}{P} \cdot \ln \frac{P_i}{P} \quad (1)$$


$\mathcal{E} = 0$ means that all the population is in one cell whereas $\mathcal{E} = 0$ means that the population is uniformly distributed.

- 3 Spatial-autocorrelation given by Moran index, with simple spatial weights given by $w_{ij} = 1/d_{ij}$


$$I = M \cdot \frac{\sum_{ij} w_{ij} (P_i - \bar{P}) \cdot (P_j - \bar{P})}{\sum_{ij} w_{ij} \sum_i (P_i - \bar{P})^2}$$

- 4 Mean distance between individuals

$$\bar{d} = \frac{1}{d_M} \cdot \sum_{i < j} \frac{P_i P_j}{P^2} \cdot d_{ij}$$




Phase transitions of indicators unveiled by exploration of the parameter space (80000 parameter points, 10 repetitions each)




Brute force calibration by exploring the parameter space. Reproduction of most existing configuration in the morphological sense (here in principal plan).


- Application on worldwide urban areas using the GHSL population grid
- Dynamical calibration on three successive time windows (1975-1990, 1990-2000, 2000-2015)
- Extraction of neighborhood (spatial span times 1.5) of 1000 larger urban areas (2015 population)
- Computation of morphological indicators on corresponding population grids




Regional patterns for Moran index





Complementarity of average distance indicator



Disjoint distributions and temporal trajectories: model application is relevant





- For a time period $[t_0, t_f]$, model initialized with population at t_0
- Growth rate N_G is computed by constraining with actual population growth: $N_G = (P_i(t_f) - P_i(t_0)) / T$ where T is a calibration parameter
- Fitness functions: squared relative population error and squared relative morphological indicator error
- Biobjective genetic algorithm (NSGA2) run for each areas (~ 600 areas large enough) and each time windows (1000 generations, population of 100), on parameters α, β, T

Performance constraints: large number of calibration algorithms to be run

→ model implemented in `scala` and integrated within the `spatialdata` library for spatial sensitivity analysis (including implementations of [Raimbault, 2018b] [Raimbault, 2018a] [Favaro and Pumain, 2011] [Cottineau, 2018])
See <https://github.com/openmole/spatialdata>



→ integration into the OpenMOLE model exploration open source software [Reuillon et al., 2013]





Enables seamlessly (i) model embedding; (ii) access to HPC resources; (iii) exploration and optimization algorithms


Apply for the OpenMOLE summer school before January 15th

<https://exmodelo.org/>




Different performance on indicator fit and forms of Pareto fronts





Urban areas trajectories can be projected into model phase diagrams



Slight significant correlations with growth rates of GHG emissions: more aggregation correlates with decreasing emissions e.g.

Implications

- This rather simple model can effectively be calibrated on worldwide urban areas and provide comparable growth parameters
- Link between urban form, urban dynamics and sustainability properties

Developments

- Benchmark with other simple models (e.g. [Li et al., 2019] presented yesterday)
- Coupling with macro-scale model for system of cities [Raimbault, 2019a]
- Investigate the link between spatial non-stationarity and non-ergodicity through simulation by the model

Within the same stream of modeling (complementary to more classical approaches to land-use change):

More realistic models?

→ Introducing more concrete ontologies, economic processes

[Bonin and Hubert, 2014], qualitative differentiation [Bonin and Hubert, 2012], governance processes [Le Néchet and Raimbault, 2015]

→ Possible bridges with Land-use change models/Land-use Transport models [Wegener and Fürst, 2004], with systems of cities models [Pumain and Reuillon, 2017]

More data-driven models?

→ More systematic link with sustainability indicators: GHG emissions, economics, etc. [Raimbault, 2019b]

→ Study models on hybrid synthetic data [Raimbault et al., 2019]: systematic conclusions for policies

→ A simple model of urban morphogenesis at the mesoscopic scale systematically calibrated: **need for more benchmarking and comparison of models.**

→ At the macro scale of the system of cities? **Need for multi-scale models.**


→ With more refined urban characteristics and other dimensions ? **Need for more interdisciplinarity.**

Use and contribute to OpenMOLE: <https://github.com/openmole>

Apply to the summer school: <https://exmodelo.org/>

- Code, data and results available at <https://github.com/JusteRaimbault/UrbanGrowth>
- Acknowledgments: Thanks to the *European Grid Infrastructure* and its *National Grid Initiatives* (*France-Grilles* in particular) to give the technical support and the infrastructure.

Reserve Slides



Examples of generated territorial shapes





Illustration of path-dependence in a simplified one-dimensional version of the model: cell trajectories in time for 9 independent repetitions from the same initial configuration.






Computation of morphological indicators on population density data for Europe (shown here on France), morphological classification.

The one-dimensional model verifies the PDE :


$$\delta t \cdot \frac{\partial p}{\partial t} = \frac{N_G \cdot p^\alpha}{P_{\alpha t}} + \frac{\alpha \beta (\alpha - 1) \delta x^2}{2} \cdot \frac{N_G \cdot p^{\alpha-2}}{P_\alpha(t)} \cdot \left(\frac{\partial p}{\partial x} \right)^2 + \frac{\beta \delta x^2}{2} \cdot \frac{\partial^2 p}{\partial x^2} \cdot \left[1 + \alpha \frac{N_G p^{\alpha-1}}{P_{\alpha t}} \right] \quad (2)$$



Stationary behavior of 1D model







Large number of repetitions show good convergence properties





- Eurostat population density raster (100m, simplified at 500m resolution)
- Overlapping (10km offset) squares of 50km side : equivalent to smoothing, removes window shape effect. Not very sensitive to window size (tested with 30km and 100km)
- Indicators computed using Fast Fourier Transform Convolution
- Classification using repeated k-means ; number of clusters taken at transition in clustering coefficient.





Potentialities of targeted model explorations: here feasible space using Pattern Space Exploration algorithm [Chérel et al., 2015].

Proposition of an interdisciplinary definition




Meta-epistemological framework of imbricated notions:




Self-organization \supseteq Morphogenesis \supseteq Autopoiesis \supseteq Life



Properties:




- Architecture links form and function
- Emergence strength [Bedau, 2002] increases with notion depth, as bifurcations [Thom, 1974]




Definition of Morphogenesis : *Emergence of the form and the function in a strongly coupled manner, producing an emergent architecture [Doursat et al., 2012]*





-  Batty, M. (1991).
Generating urban forms from diffusive growth.
Environment and Planning A, 23(4):511–544.
-  Bedau, M. (2002).
Downward causation and the autonomy of weak emergence.
Principia, 6(1):5.
-  Bonin, O. and Hubert, J.-P. (2012).
Modèle de morphogénèse urbaine: simulation d'espaces
qualitativement différenciés dans le cadre du modèle de l'économie
urbaine.




-  Bonin, O. and Hubert, J.-P. (2014).
Modélisation morphogénétique de moyen terme des villes: une schématisation du modèle théorique de ritchot et desmarais dans le cadre du modèle standard de l'économie urbaine.
Revue dEconomie Regionale Urbaine, (3):471–497.
-  Chérel, G., Cottineau, C., and Reuillon, R. (2015).
Beyond corroboration: Strengthening model validation by looking for unexpected patterns.
PLoS ONE, 10(9):e0138212.
-  Clarke, K. C., Gazulis, N., Dietzel, C., and Goldstein, N. C. (2007).
A decade of sleuthing: Lessons learned from applications of a cellular automaton land use change model.
Classics in IJGIS: twenty years of the international journal of geographical information science and systems, pages 413–427.




-  Cottineau, C., Reuillon, R., Chapron, P., Rey-Coyrehourcq, S., and Pumain, D. (2015).
A modular modelling framework for hypotheses testing in the simulation of urbanisation.
Systems, 3(4):348–377.
-  Dodds, P. S., Dewhurst, D. R., Hazlehurst, F. F., Van Oort, C. M., Mitchell, L., Reagan, A. J., Williams, J. R., and Danforth, C. M. (2017).
Simon's fundamental rich-get-richer model entails a dominant first-mover advantage.
Physical Review E, 95(5):052301.

-  Doursat, R., Sayama, H., and Michel, O. (2012).
Morphogenetic engineering: toward programmable complex systems.
Springer.
-  Engelen, G. and White, R. (2008).
Validating and calibrating integrated cellular automata based models of land use change.
In *The dynamics of complex urban systems*, pages 185–211.
Springer.
-  Favaro, J.-M. and Pumain, D. (2011).
Gibrat revisited: An urban growth model incorporating spatial interaction and innovation cycles.
Geographical Analysis, 43(3):261–286.

-  Fujita, M. and Thisse, J.-F. (1996).
Economics of agglomeration.
Journal of the Japanese and international economies,
10(4):339–378.
-  Le Néchet, F. and Raimbault, J. (2015).
Modeling the emergence of metropolitan transport authority in a
polycentric urban region.
-  Li, Y., Rybski, D., and Kropp, J. P. (2019).
Singularity cities.
Environment and Planning B: Urban Analytics and City Science,
page 2399808319843534.

-  Pumain, D. and Reuillon, R. (2017).
Urban Dynamics and Simulation Models.
Springer International.
-  Raimbault, J. (2018a).
Calibration of a density-based model of urban morphogenesis.
PloS one, 13(9):e0203516.
-  Raimbault, J. (2018b).
Indirect evidence of network effects in a system of cities.
Environment and Planning B: Urban Analytics and City Science,
page 2399808318774335.
-  Raimbault, J. (2019a).
A multi-scalar model for system of cities.
In *Conference on Complex Systems 2019*, Singapore, Singapore.

-  Raimbault, J. (2019b).
Multi-dimensional urban network percolation.
Journal of Interdisciplinary Methodologies and Issues in Science,
5:5.
-  Raimbault, J., Cottineau, C., Texier, M. L., Néchet, F. L., and
Reuillon, R. (2019).
Space matters: extending sensitivity analysis to initial spatial
conditions in geosimulation models.
Journal of Artificial Societies and Social Simulation, 22(4):10.
-  Reuillon, R., Leclaire, M., and Rey-Coyrehourcq, S. (2013).
Openmole, a workflow engine specifically tailored for the distributed
exploration of simulation models.
Future Generation Computer Systems, 29(8):1981–1990.

-  Thom, R. (1974).
Stabilité structurelle et morphogénèse.
Poetics, 3(2):7–19.
-  Turing, A. M. (1952).
The chemical basis of morphogenesis.
Philosophical Transactions of the Royal Society of London B: Biological Sciences, 237(641):37–72.
-  Wegener, M. and Fürst, F. (2004).
Land-use transport interaction: state of the art.
Available at SSRN 1434678.