

# Multiplier Stabilization Applied to Two-Stage Stochastic Programs

Clara Lage, Claudia Sagastizábal, Mikhail Solodov

► **To cite this version:**

Clara Lage, Claudia Sagastizábal, Mikhail Solodov. Multiplier Stabilization Applied to Two-Stage Stochastic Programs. 2020, pp.158-178. halshs-02900862

**HAL Id: halshs-02900862**

**<https://halshs.archives-ouvertes.fr/halshs-02900862>**

Submitted on 16 Jul 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

CES

Centre d'Économie de la Sorbonne  
UMR 8174

**Multiplier Stabilization Applied to  
Two-Stage Stochastic Programs**

Clara LAGE, Claudia SAGASTIZÁBAL, Mikhail SOLODOV

2020.10



# Multiplier Stabilization Applied to Two-Stage Stochastic Programs

Clara Lage · Claudia Sagastizábal ·  
Mikhail Solodov

Received: November 2018; Revised: April 2019

**Abstract** In many mathematical optimization applications dual variables are an important output of the solving process, due to their role as price signals. When dual solutions are not unique, different solvers or different computers, even different runs in the same computer if the problem is stochastic, often end up with different optimal multipliers. From the perspective of a decision maker, this variability makes the price signals less reliable and, hence, less useful. We address this issue for a particular family of linear and quadratic programs by proposing a solution procedure that, among all possible optimal multipliers, systematically yields the one with the smallest norm. The approach, based on penalization techniques of nonlinear programming, amounts to a regularization in the dual of the original problem. As the penalty parameter tends to zero, convergence of the primal sequence and, more critically, of the dual is shown under natural assumptions. The methodology is illustrated on a battery of two-stage stochastic linear programs.

**Keywords** Multiplier stability · Dual regularization · Penalty method · Stochastic programming · Two-stage stochastic programming · Empirical approximations

**Mathematics Subject Classification (2000)** 90C15 · 65K05 · 90C25 · 65K10 · 46N10

---

Clara Lage

ENGIE, 1, place Samuel de Champlain, 92930 Paris La Défense, France. Email: clara.lage@engie.com. Also IMPA, Rio de Janeiro, Brazil.

Claudia Sagastizábal (corresponding author)

IMECC - UNICAMP, Rua Sergio Buarque de Holanda, 651, 13083-859, Campinas, SP, Brazil. Email: sagastiz@unicamp.br. Adjunct Researcher.

Mikhail Solodov

IMPA – Instituto de Matemática Pura e Aplicada, Estrada Dona Castorina 110, Jardim Botânico, Rio de Janeiro, RJ 22460-320, Brazil. Email: solodov@impa.br.

## 1 Introduction and Motivation

Parametric optimality studies how the solution set of a nonlinear programming problem (NLP) behaves, when subject to perturbations in the objective function and/or in the constraints. For the latter, it is possible to identify circumstances in which solutions vary in a Lipschitzian-like manner with respect to right-hand side perturbations.

Studies of this type are mostly concerned with optimal values and with *primal* solution sets. In this work, we are more interested in understanding how the *dual* solutions to certain NLPs behave, when the constraints' right-hand side varies. Our motivation stems from the fact that Lagrange multipliers give the rates of change of the optimal value with respect to such perturbations. Each multiplier signals the marginal effect of raising or lowering the value of the corresponding constraint.

The interpretation of Lagrange multipliers as marginal costs (or shadow prices in the parlance of Linear Programming) has plenty of useful applications. When the Lagrange multiplier vector is not unique, the price components are related to rates of change of the subderivatives of the optimal value function. If there is a full set of prices attached to certain perturbation parameter, it is natural to ask the following important question:

*Is it possible to devise a solution methodology that provides the minimal-norm multiplier?*

The economic interest of this question is clear, since the mechanism would systematically yield the price with smallest possible norm. We provide an answer to this question for a particular family of linear and quadratic programs. To define and justify a solution procedure that, among all possible optimal multipliers, provides the smallest one in norm, we combine some variational analysis considerations and penalization techniques of nonlinear programming.

The theory is developed in the context of quadratic programming, with an application to two-stage stochastic linear programming problems, with uncertainty in the feasible set in the primal formulation. The specific problem under consideration is described in Section 4 below. In our modelling paradigm, decisions are taken independently of future observations, on the basis of the uncertainty realization, which reveals all at once; see [1, Chapter 2]. The continuous probability distribution of the stochastic variable is approximated by a finite number of Monte-Carlo scenarios, yielding finitely discrete measures, as in the sample average approximation approach [2]. In the setting of  $\mathbf{L}^p$ -spaces, a general duality approach for two-stage stochastic programs can be found in [3, 4].

Two-stage formulations are widely used; nevertheless, the paradigm remains computationally challenging in applications and has given rise to a vast literature on dedicated numerical solvers, most notably related with decomposition methods [6–8], and more recently [9–12].

Two-stage models are well suited to situations in which the output of interest is the first-stage solution, a deterministic value decided “here-and-now”. Another useful output is related to (some components) of the multiplier in

the affine constraint. This is a stochastic variable of the “wait-and-see” type, whose expected value gives a price signal that helps defining business strategies. Having this goal in mind, the fact that different samples can make the price signal vary wildly is a serious handicap. Our proposal addresses this issue, by systematically providing the minimal-norm price signal, thus making the indicator more reliable for the decision maker.

The rest of the paper is organized as follows. In Section 2, we support with some Variational Analysis considerations our computational approach to building multiplier estimates converging to minimal-norm optimal ones. We also introduce an apparently new condition, which allows to prove boundedness of the set of Lagrange multipliers associated with some part of the constraints of the problem, while allowing the other multipliers to be unbounded. In Section 3, we prove convergence of the proposed multiplier estimates (obtained from an exterior penalty scheme) to the ones of minimal norm, for linear or quadratic programs satisfying some natural assumptions. In Section 4, we explain how the methodology can be applied to two-stage stochastic linear programming problems, and present numerical results that confirm the interest of the approach. The work ends with some concluding remarks.

## 2 A Variational Analysis Perspective

We mostly follow the notation of [13]. Points in  $\mathbb{R}^n$  are considered as column vectors. The Euclidean inner product and norm are denoted by  $\langle \cdot, \cdot \rangle$  and  $\| \cdot \|$ , respectively. The indicator function of a set  $S$  is denoted by  $\delta_S(\cdot)$ , i.e., this function is 0 for points in  $S$  and is  $+\infty$  otherwise. If  $S$  is convex, then  $N_S(x)$  stands for the normal cone of  $S$  at the point  $x$ . The unit ball centered at 0 is  $\mathbb{B}$ , and the identity matrix is  $I$ ; in both cases the dimension is always clear from the context. For a proper convex function  $f$ , its subdifferential at  $x$  is denoted by  $\partial f(x)$ , while its horizon subdifferential at the point  $x$  is the normal cone of the function’s domain, i.e.,  $\partial^\infty f(x) = N_{\text{dom } f}(x)$ ; see [13, Proposition 8.12].

We are particularly interested in the Lagrange multiplier of the affine equality constraint of the following (feasible) optimization problem:

$$\min f(x) \text{ s.t. } x \in X, Ax - b = 0, \quad (1)$$

where  $f$  is a finite-valued convex and continuously differentiable function,  $X$  is a closed convex set, and  $b \in \{y : y = Ax, x \in X\}$ .

While this is not essential for some of the subsequent considerations, we shall assume that  $X$  is defined by smooth convex inequalities, as is certainly the case in the applications we have in mind. It is then well known that uniqueness of Lagrange multipliers associated to a solution  $\bar{x}$  of problem (1) is implied by the linear independence of gradients of the constraints active at  $\bar{x}$ . This is in turn equivalent to the so-called strict Mangasarian-Fromovitz (MF) condition (see [14] and also [15, Sections 1.1, 1.2.4]). We emphasize that in (1) either of these assumptions implies that the matrix  $A$  is of full rank, a condition that rarely holds in practice for many important applications. The less stringent MF

constraint qualification (MFCQ), equivalent to having a nonempty compact set of Lagrange multipliers, also subsumes that  $A$  has full rank.

Thus, if  $A$  is not of full rank, the multipliers associated to the equality constraint in (1) are necessarily not unique. In fact, since MFCQ is violated in this case, the multiplier set is unbounded. This leads us to focus on devising a mechanism to identify/compute the multiplier that has the minimal norm. The idea is to consider a sequence of problems that penalize the equality constraint in (1), depending on a parameter  $\beta > 0$ . Given a (primal) solution to the penalized problem, we then construct an explicit multiplier estimate, which we denote by  $\pi^\beta$ . Specifically, we solve

$$\min f(x) + \frac{1}{2\beta} \|Ax - b\|^2 \quad \text{s.t. } x \in X,$$

for  $\beta > 0$  to obtain  $x^\beta$ , and define as multiplier proxy

$$\pi^\beta := \frac{Ax^\beta - b}{\beta}.$$

For the case when (1) is a linear or quadratic program, including the two-stage stochastic linear programming problems considered in Section 4, we then exhibit some natural conditions, which ensure that, as  $\beta \rightarrow 0$ , the sequence of the constructed multiplier estimates  $\pi^\beta$  tends to the specific multiplier  $\hat{\pi}$  of minimal norm. The precise details will be given in Section 3.

Approximating Lagrange multipliers in the setting of quadratic penalty methods, along the lines above, is certainly not a new idea; see, e.g., [16, Chapter 17.1]. However, in the literature convergence results are established assuming linear independence of active gradients (as well as subsequential convergence of the primal sequence  $x^\beta$ ), in which case the optimal multiplier is unique; see [16, Theorem 17.2] and Theorem 3.2 below. In Section 3, we give conditions under which  $x^\beta$  converges, and show convergence of  $\pi^\beta$  (to minimal-norm multiplier), without assuming the linear independence condition, thus covering a much more general case.

In this section, we give a different motivation and insight for the multiplier proxies  $\pi^\beta$ , by specializing some results of Variational Analysis [13] to our setting. We start with a fixed  $\beta \geq 0$ , and relate the estimates  $\pi^\beta$  with a particular instance of the *generalized Lagrange multiplier* rule [13, Example 10.8, p.429]. More precisely, given a scalar  $\beta \geq 0$ , consider the following penalties:

$$\mathbb{R}^m \ni v \mapsto \theta^\beta(v) := \sup_{y \in \mathbb{R}^m} \left\{ \langle v, y \rangle - \frac{1}{2}\beta \|y\|^2 \right\} = \begin{cases} \frac{1}{2\beta} \|v\|^2 & \text{if } \beta > 0 \\ \delta_{\{0\}}(v) & \text{if } \beta = 0. \end{cases} \quad (2)$$

These (lower-semicontinuous, proper, convex) functions are a particular case of the piecewise linear-quadratic penalties in [13, Example 11.18, p. 497] (therein,  $\theta^\beta$  corresponds to  $\theta_{Y,B}$ , written for  $Y = \mathbb{R}^m$  and the, possibly null, matrix  $B = \beta I$ ). The respective subdifferentials are:

$$\begin{aligned} & \text{if } \beta > 0, \quad \text{for all } v \in \mathbb{R}^m = \text{dom } \theta^\beta, \\ & \partial \theta^\beta(v) = \left\{ \frac{1}{\beta} v \right\} \quad \text{and} \quad \partial^\infty \theta^\beta(v) = \{0\}, \end{aligned} \quad (3)$$

while

$$\text{if } \beta = 0, \quad \text{for } v = 0 = \text{dom } \theta^0, \quad \partial\theta^0(v) = \mathbb{R}^m \text{ and } \partial^\infty\theta^0(v) = \mathbb{R}^m. \quad (4)$$

The connection between penalties and dual variables (multipliers) is made clear when considering, for perturbation parameters  $u \in \mathbb{R}^m$ , the (unconstrained) parametric minimization problems

$$\min_{x \in \mathbb{R}^n} f^\beta(x, u) := f(x) + \delta_X(x) + \theta^\beta(Ax - b + u), \quad (5)$$

noting that writing (5) with  $\beta = 0$  and  $u = 0$  yields our original problem (1).

When  $\beta > 0$ , some  $x^\beta$  is optimal in (5) if and only if

$$x^\beta \in X, \quad \mu^\beta \in N_X(x^\beta), \quad \nabla f(x^\beta) + \mu^\beta + A^\top \pi^\beta = 0, \quad (6)$$

where, for  $\bar{u} \in \mathbb{R}^m$  given, the unique *extended Lagrange multiplier* in [13] is

$$\pi^\beta := \frac{Ax^\beta - b + \bar{u}}{\beta}.$$

To consider the case when  $\beta = 0$ , recall that, in its dual formulation (see [14] and also [15, Sections 1.1, 1.2.4]), the MFCQ at a feasible point  $\bar{x}$  of our (unperturbed) problem (1) means that

$$0 = A^\top \pi + \mu, \quad \mu \in N_X(\bar{x}) \quad \Rightarrow \quad \pi = 0 \in \mathbb{R}^m \text{ and } \mu = 0 \in \mathbb{R}^n. \quad (7)$$

If  $\bar{x}$  satisfies MFCQ (7), there exists a classical Lagrange multiplier  $\bar{\pi} \in \mathbb{R}^m$ , not necessarily unique, satisfying (6), written with  $\beta = 0$  and  $(x^\beta, \pi^\beta, \mu^\beta) = (\bar{x}, \bar{\pi}, \bar{\mu})$ . Condition (6) is also sufficient for  $\bar{x}$  to be optimal for (5), written with  $\beta = 0$  and  $\bar{u} := b - A\bar{x}$ .

The proposition below, that follows from the *parametric version of Fermat rule* in [13, Example 10.12], analyzes (5) from a Variational Analysis perspective, condensing the key ingredients relating extended Lagrange multipliers to the marginal rate of change of the optimal value in (1), when considered as a function of the right-hand side perturbation of the affine constraint.

**Proposition 2.1 (Extended Lagrange multipliers)** *Associated to (1), consider the parametric optimization problems (5) with penalties (2), where  $u = \bar{u} \in \mathbb{R}^m$  and  $\beta \geq 0$  are fixed. Let the corresponding optimal value and solution set be given by*

$$p^\beta(u) := \inf_{x \in \mathbb{R}^n} f^\beta(x, u) \quad \text{and} \quad P^\beta(u) := \arg \min_{x \in \mathbb{R}^n} f^\beta(x, u).$$

The following holds.

- (i) When  $\beta > 0$ , the function  $p^\beta$  is convex, strictly differentiable at any point  $\bar{u} \in \text{dom } p^\beta = \mathbb{R}^m$ , with gradient  $\nabla p^\beta(\bar{u}) = \pi^\beta$ .
- (ii) When  $\beta = 0$ , the function  $p^0$  is convex, strictly continuous at the point  $\bar{u} = A\bar{x} - b = \text{dom } p^0$ , with subdifferential

$$\partial p^0(\bar{u}) = \{\bar{\pi} \in \mathbb{R}^m : (6) \text{ holds written with } (x^\beta, \pi^\beta, \mu^\beta) = (\bar{x}, \bar{\pi}, \bar{\mu})\}. \quad (8)$$

**Proof** The perturbed function  $f^\beta(x, u)$  is convex in  $(x, u)$ , and the finite-valued function  $f$  has full domain. In this situation, by [13, Example 10.8],

$$\partial f^\beta(\bar{x}, \bar{u}) = \nabla f(\bar{x}) + N_X(\bar{x}) + \partial \theta^\beta(A\bar{x} - \bar{u}), \quad \partial^\infty f^\beta(\bar{x}, \bar{u}) = \partial^\infty \theta^\beta(A\bar{x} - \bar{u}).$$

Since  $f^\beta$  is convex (therefore regular) with our definitions, the  $Y$ -sets in [13, Theorem 10.13] satisfy the relations

$$Y(\bar{u}) = \{ \pi : (0, \pi) \in \partial f^\beta(x^\beta, \bar{u}) \} \text{ and } Y^\infty(\bar{u}) = \{ \pi : (0, \pi) \in \partial^\infty f^\beta(x^\beta, \bar{u}) \},$$

for any  $x^\beta \in P^\beta(\bar{u})$  and  $\bar{u} \in \text{dom } p^\beta = \text{dom } \theta^\beta$ . Together with (3) and (4), this gives  $\partial^\infty p^\beta(\bar{u}) = Y^\infty(\bar{u}) = \partial^\infty \theta^\beta(Ax^\beta - \bar{u})$  and, as claimed,

$$\partial p^\beta(\bar{u}) = Y(\bar{u}) = \{ \pi^\beta : (x^\beta, \mu^\beta, \pi^\beta) \text{ satisfies (6)} \}. \quad \square$$

The above characterization, obtained from the penalty scheme as extended Lagrange multiplier, motivates from the Variational Analysis point of view the choice of the multiplier estimates in our development. In Section 3, we shall show under which conditions such estimates converge to the minimal-norm multipliers. Among other things, we shall need the following result, which establishes boundedness of the set of Lagrange multipliers associated to some part of the constraints of the problem, while allowing the other multipliers to be unbounded. Apparently, this result is new.

Recall the MFCQ condition (7) is equivalent to the set of multipliers being nonempty and bounded. Consider the following condition at  $\bar{x}$  feasible in (1):

$$0 = A^\top \pi + \mu, \quad \mu \in N_X(\bar{x}) \quad \Rightarrow \quad \mu = 0. \quad (9)$$

Clearly, (9) is a weaker condition than (7). In particular, as we show in Theorem 2.1 below, (9) implies boundedness only for the  $\mu$ -part of the multipliers, while the  $\pi$ -part can be unbounded. The condition in question can be interpreted as a ‘‘partial’’ MFCQ condition. However, note that (9) is *not* a constraint qualification, i.e., it does not imply (by itself) that for a solution  $\bar{x}$  of problem (1) the multiplier set is nonempty. An alternative, equivalent, formulation of condition (9) is

$$\text{Im } A^\top \cap N_X(\bar{x}) = \{0\}. \quad (10)$$

**Theorem 2.1 (On boundedness of multipliers)** *Let  $\bar{x}$  be any feasible point in (1). Then the following statements are equivalent:*

- (i) *Condition (9) holds at  $\bar{x}$ .*
- (ii) *For any  $\bar{g} \in \mathbb{R}^n$ , the set*

$$S_{\bar{g}} := \{ \bar{\mu} \in N_X(\bar{x}) : \exists \bar{\pi} \in \mathbb{R}^m \text{ s.t. } \bar{g} + A^\top \bar{\pi} + \bar{\mu} = 0 \}$$

*is bounded.*



**Proof** We shall show the equivalent assertion

$$\exists \tilde{\mu} \in \text{Im } A^\top \cap N_X(\bar{x}), \tilde{\mu} \neq 0 \iff \exists \bar{g} \in \mathbb{R}^n \text{ such that } S_{\bar{g}} \text{ is unbounded.}$$

Assume first that for some  $\bar{g}$  the set  $S_{\bar{g}}$  is unbounded, i.e., there exists a sequence  $\{(\pi^k, \mu^k)\}$  such that

$$\bar{g} + A^\top \pi^k + \mu^k = 0, \quad \mu^k \in N_X(\bar{x}), \quad (11)$$

with  $\|\mu^k\| \rightarrow +\infty$ . As  $N_X(\bar{x})$  is a closed cone, we can assume, passing onto a subsequence if necessary, that

$$\mu^k / \|\mu^k\| \rightarrow \bar{\mu} \in N_X(\bar{x}), \quad \bar{\mu} \neq 0.$$

Denote  $u^k = -A^\top \pi^k / \|\mu^k\| \in \text{Im } A^\top$ . Dividing the equality in (11) by  $\|\mu^k\|$  and passing onto the limit, it follows that

$$u^k = (\bar{g} + \mu^k) / \|\mu^k\| \rightarrow \bar{\mu}.$$

As  $u^k \in \text{Im } A^\top$ ,  $u^k \rightarrow \bar{\mu}$ , and  $\text{Im } A^\top$  is closed, we conclude that  $\bar{\mu} \in \text{Im } A^\top$ . As it also holds that  $\bar{\mu} \in N_X(\bar{x})$  and  $\bar{\mu} \neq 0$ , this contradicts (9).

Suppose now that there exists  $0 \neq \tilde{\mu} \in N_X(\bar{x})$  such that  $A^\top \tilde{\pi} + \tilde{\mu} = 0$  for some  $\tilde{\pi}$ . If for some  $\bar{g}$  there is a pair  $(\bar{\pi}, \bar{\mu})$  satisfying

$$\bar{g} + A^\top \bar{\pi} + \bar{\mu} = 0, \quad \bar{\mu} \in N_X(\bar{x}),$$

then, for any  $t > 0$ , it holds that  $\bar{\mu} + t\tilde{\mu} \in N_X(\bar{x}) + N_X(\bar{x}) = N_X(\bar{x})$ , since the cone in question is convex. Hence, for any  $t > 0$ ,

$$\bar{g} + A^\top (\bar{\pi} + t\tilde{\pi}) + (\bar{\mu} + t\tilde{\mu}) = 0, \quad \bar{\mu} + t\tilde{\mu} \in N_X(\bar{x}).$$

Since  $\tilde{\mu} \neq 0$ , as  $t \rightarrow +\infty$ ,  $\|\bar{\mu} + t\tilde{\mu}\| \rightarrow +\infty$ , and the set  $S_{\bar{g}}$  is unbounded.  $\square$

We emphasize that, being weaker than MFCQ, condition (9) is certainly not restrictive (assuming that the existence of Lagrange multipliers is given or follows from some other considerations).

### 3 A Nonlinear Programming Computational Perspective

Consider now the following (linear or) *quadratic programming* problem:

$$\min f(x) := \langle g, x \rangle + \frac{1}{2} \langle x, Hx \rangle \quad \text{s.t. } x \in X, Ax - b = 0, \quad (12)$$

where  $X := \{x \in \mathbb{R}^n : x \geq 0\}$  and  $b \in \{y : y = Ax, x \in X\}$ ,  $g \in \mathbb{R}^n$  and  $H$  is an  $n \times n$  symmetric matrix ( $H = 0$  corresponding to linear programming). We note that in our developments below,  $H$  is not necessarily positive semidefinite, although it also might be.

When  $H$  is positive semidefinite, the convex problem (12) is a particular instance of (1), and we can use the constructions in Section 2 for some motivations. In that case, fixing  $\bar{u} = 0$ , for  $\beta > 0$  from (2) and (5), we have

$$f^\beta(x, 0) = f(x) + \delta_X(x) + \frac{1}{2\beta} \|Ax - b\|^2,$$

and Proposition 2.1 characterizes the extended Lagrange multiplier as follows:

$$\pi^\beta = \frac{Ax^\beta - b}{\beta}, \quad \text{for } x^\beta \in P^\beta(0).$$

As a result, finding  $x^\beta \in P^\beta(0)$  is equivalent to finding  $x^\beta$ , a solution to the following (partial) exterior penalization of problem (12):

$$\min f(x) + \frac{1}{2\beta} \|Ax - b\|^2 \quad \text{s.t. } x \in X. \quad (13)$$

The multiplier estimate is then given by

$$\pi^\beta := \frac{Ax^\beta - b}{\beta}. \quad (14)$$

Penalty methods (see, e.g., [17]) solve subproblems (13) for a sequence of decreasing penalty parameters  $0 < \beta_{k+1} < \beta_k$ , tending to zero. We want to study how the multiplier estimates (14) for the equality constraints in (12) behave along the sequence of solving the penalized subproblems (13). We shall show that, under reasonable assumptions,  $\pi^\beta$  converge to the minimal-norm multiplier  $\hat{\pi}$ ; see (24) below for a formal definition.

We start with some standard facts on (primal) convergence of penalty methods [17], that do not depend on the setting of (12), and can also use other forms of exterior penalties (not necessarily quadratic). But we shall keep this setting for the sake of not introducing extra notation. Define

$$F_k(x) := f(x) + \frac{1}{2\beta_k} \|Ax - b\|^2,$$

the objective function in (13).

**Theorem 3.1 (Primal convergence of generic penalty methods)** *Let  $x^k$  be a (global) solution of (13) for  $\beta = \beta_k$  for each  $k$ , with  $0 < \beta_{k+1} < \beta_k$ . Then,*

$$F_{k+1}(x^{k+1}) \geq F_k(x^k), \quad \|Ax^{k+1} - b\| \leq \|Ax^k - b\|, \quad f(x^{k+1}) \geq f(x^k). \quad (15)$$

*If, in addition,  $\beta_k \rightarrow 0$  as  $k \rightarrow \infty$ , and the optimal value of problem (12) is finite, then every accumulation point of  $\{x^k\}$  is a (global) solution of (12).*

Note that this result refers to *global* solutions of subproblems. This is standard, and also not an issue when the problem (12) is convex. Another observation is that in the case of a quadratic program as ours, if  $f$  is bounded below on the feasible region (i.e., the optimal value is finite), then problem (12) has a solution, by the Frank–Wolfe Theorem [18].

However, it is important to emphasize that the general convergence result in Theorem 3.1 asserts optimality of accumulation points, but does not say anything about their *existence*. It can thus be “vacuous”, if the sequence is unbounded. Our first task will be to prove when the generated sequence  $\{x^k\}$  is bounded. But before proceeding, we shall mention the following classical result on convergence of the multiplier estimates, obtained from the quadratic penalty method. Let  $x^k$  be a solution of (13) for  $\beta = \beta_k$ . Define

$$\pi^k := \frac{1}{\beta_k}(Ax^k - b). \quad (16)$$

The assertion below is standard; see, e.g., [16, Theorem 17.2]. Like Theorem 3.1 above, it does not depend on the setting of problem (12), and can be easily extended to the case of general nonlinear objective function  $f$  and general nonlinear constraints, including inequality constraints. As this is not essential for our developments, we state the result for equality constraints only.

**Theorem 3.2 (Dual convergence of the quadratic penalty method)**

*In (12), let  $X = \mathbb{R}^n$ . Let  $\bar{x}$  be any accumulation point of  $\{x^k\}$ , where  $x^k$  is a solution of (13) for  $\beta = \beta_k$  for each  $k$ ,  $\{x^{k_j}\} \rightarrow \bar{x}$  as  $j \rightarrow \infty$ . Let the linear independence constraints qualification hold at  $\bar{x}$  (in the setting of (12) this means that  $A$  has full rank).*

*Then,  $\bar{x}$  is a stationary point of (12) and the subsequence  $\{\pi^{k_j}\}$ , defined by (16), converges to the unique Lagrange multiplier  $\bar{\pi}$  associated to  $\bar{x}$ .*

Note, however, that Theorem 3.2 again implicitly assumes boundedness of  $\{x^k\}$  (as it refers to its accumulation points), and requires the linear independence constraints qualification for convergence of the dual sequence. The latter, in particular, is not assumed in our setting.

For establishing boundedness of the primal sequence, we shall need the following conditions. Recall that the standard critical cone of (12) at a given stationary point  $\bar{x}$  is defined by

$$K(\bar{x}) := \text{Ker } A \cap \{d \in \mathbb{R}^n : \langle H\bar{x} + g, d \rangle \leq 0, d_i \geq 0 \text{ for } i \text{ s.t. } \bar{x}_i = 0\}. \quad (17)$$

In the case at consideration, the Hessian of the Lagrangian (for any point  $(\bar{x}, \bar{\pi}, \bar{\mu})$ ) is the matrix  $H$ . Thus, the usual second-order sufficient optimality condition for  $\bar{x}$  states that

$$\langle Hd, d \rangle > 0 \quad \text{for all } d \in K(\bar{x}) \setminus \{0\}. \quad (18)$$

When  $H$  is positive semidefinite, the solution set of problem (12) is convex. Since (18) implies that  $\bar{x}$  is a strict (thus isolated) minimizer, the condition

means that in the convex case the primal solution must be unique. In particular, when  $f$  is linear, i.e.,  $H = 0$ , condition (18) holds if and only if  $K(\bar{x}) = \{0\}$ . It can be further seen that this means that  $\langle g, d \rangle > 0$  for all feasible directions  $d$  at  $\bar{x}$ . This, in turn, is equivalent to saying that  $\bar{x}$  is the unique solution of the linear program (12). Thus, for linear programming, including the two-stage stochastic linear programming setting in Section 4, the assumption (18) amounts to stating that the primal solution of the problem is unique.

Note also that since  $K(\bar{x}) \subset \text{Ker } A$ , the following is also a second-order sufficient optimality condition (as it implies (18)):

$$\langle Hd, d \rangle > 0 \quad \text{for all } d \in \text{Ker } A \setminus \{0\}. \quad (19)$$

However, unlike (18), condition (19) is an assumption on  $H$  and  $A$ , which does not depend on  $\bar{x}$ . Note that (19) does not require  $H$  to be positive semidefinite, and thus the objective function  $f$  in (12) can be non-convex.

**Theorem 3.3 (Conditions for primal convergence)** *Suppose that one of the following two items holds:*

1. *Condition (19) is satisfied.*
2. *The matrix  $H$  is positive semidefinite, and (18) holds for the solution  $\bar{x}$  of (12) (if  $H = 0$ , this amounts to (12) having a unique primal solution).*

*Then, for any sequence of parameters  $\beta_k \rightarrow 0$  (even not necessarily monotone), any sequence  $\{x^k\}$  generated by the penalty scheme (13) is bounded.*

*If also  $\beta_{k+1} < \beta_k$  for all  $k$ , then each of the accumulation points of  $\{x^k\}$  is a solution of (12). In particular, in the second case above, the whole sequence converges to the unique solution  $\bar{x}$ .*

**Proof** We reason by contradiction: taking a subsequence if necessary, suppose that  $\|x^k\| \rightarrow \infty$ .

Define  $z^k = x^k / \|x^k\|$ . Again passing onto a subsequence if necessary, we can assume that  $z^k \rightarrow z$ ,  $z \neq 0$ .

By the KKT optimality conditions for the subproblems (13), it holds that

$$Hx^k + g + \frac{1}{\beta_k} A^\top (Ax^k - b) - \mu^k = 0, \quad x^k \geq 0, \quad \mu^k \geq 0, \quad \langle \mu^k, x^k \rangle = 0. \quad (20)$$

Note that  $\langle \mu^k, z^k \rangle = 0$ . Thus, multiplying the first relation above by  $z^k$  yields

$$\langle Hx^k + g, z^k \rangle = \frac{1}{\beta_k} \langle A^\top (b - Ax^k), z^k \rangle.$$

Next, multiplying both sides of the latter equality by  $\beta_k / \|x^k\|$ , we conclude that

$$\beta_k \langle Hz^k, z^k \rangle + \frac{\beta_k}{\|x^k\|} \langle g, z^k \rangle = \frac{1}{\|x^k\|} \langle A^\top b, z^k \rangle - \|Az^k\|^2.$$

As  $\{z^k\}$  is bounded, while  $\|x^k\| \rightarrow \infty$  and  $\beta_k \rightarrow 0$ , passing onto the limit as  $k \rightarrow \infty$  yields that  $0 = \|Az\|^2$ , i.e.,  $z \in \text{Ker } A$ .

Let  $\tilde{x}$  be any feasible point in (12). Since  $\tilde{x} \in X$ ,  $A\tilde{x} - b = 0$ , and  $x^k$  is a solution of (13), it holds that

$$f(\tilde{x}) \geq f(x^k) + \frac{1}{2\beta_k} \|Ax^k - b\|^2 \geq f(x^k). \quad (21)$$

Dividing this inequality by  $\|x^k\|^2$ , we obtain that

$$\frac{f(\tilde{x})}{\|x^k\|^2} \geq \frac{f(x^k)}{\|x^k\|^2} = \frac{1}{2} \langle Hz^k, z^k \rangle + \frac{1}{\|x^k\|} \langle g, z^k \rangle.$$

Passing onto the limit as  $k \rightarrow \infty$  gives

$$0 \geq \frac{1}{2} \langle Hz, z \rangle. \quad (22)$$

Since  $0 \neq z \in \text{Ker } A$ , this immediately gives a contradiction if the condition (19) holds.

Suppose now  $H$  is positive semidefinite, and (18) holds for the solution  $\bar{x}$  of (12), which is unique in this case. Since  $\bar{x}$  is in particular feasible, from (21) written with  $\tilde{x} = \bar{x}$ , using also the convexity of  $f$ , we conclude that

$$f(\bar{x}) \geq f(x^k) \geq f(\bar{x}) + \langle \nabla f(\bar{x}), x^k - \bar{x} \rangle,$$

and, hence,

$$0 \geq \langle \nabla f(\bar{x}), x^k - \bar{x} \rangle.$$

Dividing both sides above by  $\|x^k\|$ , and passing onto the limit, we conclude that  $\langle \nabla f(\bar{x}), z \rangle \leq 0$ . Since  $z \geq 0$  is obvious (because  $x^k \geq 0$ ), and recalling that  $z \in \text{Ker } A$ , we obtain that  $0 \neq z \in K(\bar{x})$ ; see (17). Now (22) again gives a contradiction with (18). We conclude that  $\{x^k\}$  is bounded. The other assertions follow from the general results about penalty methods in Theorem 3.1 (and other considerations stated above).  $\square$

Having established when there is primal convergence of solutions of the penalized subproblems (13), we now analyze the asymptotic behavior of the dual sequence  $\{\pi^k\}$  defined by (16).

Recall that for a solution  $\bar{x}$  of problem (12), the set of associated Lagrange multipliers  $(\pi, \mu)$  is characterized by the following system:

$$H\bar{x} + g + A^\top \pi - \mu = 0, \quad \bar{x} \geq 0, \quad \mu \geq 0, \quad \langle \mu, \bar{x} \rangle = 0. \quad (23)$$

To exhibit the specific dual behavior (dual limit) of the sequence  $\{\pi^k\}$ , denote by  $\hat{\pi} = \hat{\pi}(\bar{x}, \bar{\mu})$  the minimal-norm element, which solves (23) for the given  $\bar{x}$  and  $\bar{\mu}$ , i.e., the (unique) solution of

$$\min \frac{1}{2} \|\pi\|^2 \quad \text{s.t.} \quad H\bar{x} + g + A^\top \pi - \bar{\mu} = 0. \quad (24)$$

We have the following.

**Theorem 3.4 (Convergence of the multipliers estimates)** *Let  $\beta_k \rightarrow 0$  and  $\beta_{k+1} < \beta_k$  for all  $k$ . Let the assumptions of Theorem 3.3 hold. Let  $\bar{x}$  be any accumulation point of the sequence  $\{x^k\}$  (which is bounded by Theorem 3.3),  $x^{k_j} \rightarrow \bar{x}$  as  $j \rightarrow \infty$ . Let condition (9) hold at  $\bar{x}$ .*

*Then, the sequence  $\{\mu^{k_j}\}$  is bounded. Moreover, for any of its accumulation point  $\bar{\mu}$ , the subsequence  $\{\pi^{k_j}\}$  defined by (16) converges to  $\hat{\pi}$ , the minimal-norm solution of (24). The point  $(\bar{x}, \hat{\pi}, \bar{\mu})$  is a primal-dual solution of (12).*

**Proof** Under the assumptions of Theorem 3.3, it follows that  $\{x^k\}$  is bounded. Recalling the subproblem KKT conditions (20), and using the definition (16) of  $\pi^k$ , we have that

$$Hx^k + g + A^\top \pi^k - \mu^k = 0, \quad x^k \geq 0, \quad \mu^k \geq 0, \quad \langle \mu^k, x^k \rangle = 0. \quad (25)$$

Let  $\{x^{k_j}\} \rightarrow \bar{x}$  as  $j \rightarrow \infty$ . We first prove that the sequence  $\{\mu^{k_j}\}$  is bounded. Similarly to the first part of the proof of Theorem 2.1, suppose by contradiction that (25) holds with  $\|\mu^{k_j}\| \rightarrow +\infty$  (possibly passing onto a subsequence). We can assume, passing onto a further subsequence if necessary, that

$$\mu^{k_j} / \|\mu^{k_j}\| \rightarrow \bar{\mu} \geq 0, \quad \bar{\mu} \neq 0. \quad (26)$$

Denote  $u^{k_j} = A^\top \pi^{k_j} / \|\mu^{k_j}\| \in \text{Im } A^\top$ . Dividing the equality in (25) by  $\|\mu^{k_j}\|$ , and passing onto the limit as  $j \rightarrow \infty$ , it follows that

$$u^{k_j} = (\mu^{k_j} - Hx^{k_j} - g) / \|\mu^{k_j}\| \rightarrow \bar{\mu},$$

where boundedness of  $\{x^{k_j}\}$  was taken into account. As  $u^{k_j} \in \text{Im } A^\top$ ,  $u^{k_j} \rightarrow \bar{\mu}$ , and  $\text{Im } A^\top$  is closed, we conclude that  $\bar{\mu} \in \text{Im } A^\top$ . Obviously  $\bar{x} \geq 0$  and, dividing the last two relations in (25) by  $\|\mu^{k_j}\|$  and passing onto the limit,  $\bar{\mu} \geq 0$ ,  $\langle \bar{\mu}, \bar{x} \rangle = 0$ . This means that  $-\bar{\mu} \in N_X(\bar{x})$ , where  $X = \mathbb{R}_+^n$ . As  $\bar{\mu} \neq 0$ , and  $-\bar{\mu} \in \text{Im } A^\top$ , we obtain a contradiction with (9).

Once  $\{\mu^{k_j}\}$  is bounded, the first equality in (25) implies that  $\{A^\top \pi^{k_j}\}$  is bounded as well. Passing onto a further subsequence if necessary, we can assume that  $\{x^{k_j}\} \rightarrow \bar{x}$ ,  $\{\mu^{k_j}\} \rightarrow \bar{\mu}$ ,  $\{A^\top \pi^{k_j}\} \rightarrow a$  as  $j \rightarrow \infty$ .

Taking any point  $\tilde{x}$  such that  $A\tilde{x} - b = 0$ , we observe that

$$\pi^k = \frac{1}{\beta_k}(Ax^k - b) = \frac{1}{\beta_k}A(x^k - \tilde{x}) \in \text{Im } A.$$

Thus,  $A^\top \pi^k \in \text{Im } A^\top A$ . Because this subspace is closed, we have  $a \in \text{Im } A^\top A$ , i.e.,  $a = A^\top \bar{\pi}$  for some  $\bar{\pi} \in \text{Im } A$ .

Passing onto the limit in (25) as  $j \rightarrow \infty$ , we then have that

$$H\bar{x} + g + A^\top \bar{\pi} - \bar{\mu} = 0, \quad \bar{\pi} \in \text{Im } A, \quad \bar{x} \geq 0, \quad \bar{\mu} \geq 0, \quad \langle \bar{\mu}, \bar{x} \rangle = 0. \quad (27)$$

We next show that there exists only one  $\bar{\pi} \in \text{Im } A$  which satisfies the left equality in (27), for the given  $\bar{x}$  and  $\bar{\mu}$ . Let  $\tilde{\pi}$  be any other element in  $\text{Im } A$  such that  $H\bar{x} + g + A^\top \tilde{\pi} - \bar{\mu} = 0$ . Subtracting this equality from the first one in (27), we conclude that

$$(\bar{\pi} - \tilde{\pi}) \in \text{Ker } A^\top, \quad (\bar{\pi} - \tilde{\pi}) \in \text{Im } A.$$

As  $\text{Ker } A^\top = (\text{Im } A)^\perp$ , it follows that  $\bar{\pi} = \hat{\pi}$ , i.e., the element with the properties under consideration is unique. Observe further that the solution  $\hat{\pi}$  of (24) satisfies those properties: it exists, is unique, and  $H\bar{x} + g + A^\top \hat{\pi} - \bar{\mu} = 0$ . Further, by the optimality condition for (24), it holds that there exists some  $\lambda$  such that  $\hat{\pi} + A\lambda = 0$ , i.e.,  $\hat{\pi} \in \text{Im } A$ . As we have shown that such an element is unique, it follows that  $\bar{\pi} = \hat{\pi}$ .

In particular,  $\{A^\top \pi^{k_j}\} \rightarrow a = A^\top \bar{\pi}$  now means that  $\{A^\top (\pi^{k_j} - \hat{\pi})\} \rightarrow 0$ , as  $j \rightarrow \infty$ . Finally, we show that this implies that  $\pi^{k_j} \rightarrow \hat{\pi}$  (recall that  $(\pi^{k_j} - \hat{\pi}) \in \text{Im } A$ ).

To that end, recall that, for any matrix  $A$  there exists  $\gamma > 0$  such that

$$\|A^\top Au\| \geq \gamma \|Au\| \quad \text{for all } u.$$

(To see this, assume the contrary, i.e., that there exists  $\{u^k\}$  such that  $Au^k \neq 0$ , and  $\|A^\top Au^k\|/\|Au^k\| \rightarrow 0$ . Passing onto a further subsequence, if necessary,  $Au^k/\|Au^k\| \rightarrow v \in \text{Im } A$ ,  $v \neq 0$ ,  $A^\top v = 0$ . This gives a contradiction, since  $\text{Ker } A^\top \cap \text{Im } A = \{0\}$ .)

As  $(\pi^{k_j} - \hat{\pi}) \in \text{Im } A$ , there exists some  $b^j$  such that  $\pi^{k_j} - \hat{\pi} = Ab^j$ . Then,

$$\begin{aligned} \|A^\top (\pi^{k_j} - \hat{\pi})\| &= \|A^\top Ab^j\| \geq \gamma \|Ab^j\| \\ &= \gamma \|\pi^{k_j} - \hat{\pi}\|, \end{aligned}$$

implying the assertion, since the left-hand side tends to zero as  $j \rightarrow \infty$ .  $\square$

The results presented so far provide a constructive answer to our initial question, on how to devise a solution methodology yielding the minimal-norm multiplier. The mechanism is applied in the next section to an important class of stochastic optimization problems, with linear objective function and affine constraints, and where uncertainty is dealt with by sample average approximations in two stages, via the so-called recourse functions; see (30) below.

#### 4 Application to Two-Stage Stochastic Linear Programming

For a random variable  $\xi$  with finite support, we consider two-stage stochastic linear programs with relatively complete recourse, with uncertainty in the second-stage costs  $q(\xi)$ , and the right-hand side vector  $h(\xi)$ . Specifically, the problem is

$$\min f(x) \quad \text{s.t. } x_1 \geq 0, x_2(\xi) \geq 0, Tx_1 + Wx_2(\xi) = h(\xi) \quad \text{a.e. } \xi,$$

where

$$f(x) := \langle c, x_1 \rangle + \mathbb{E}[\langle q(\xi), x_2(\xi) \rangle].$$

In this problem  $x_i \in \mathbb{R}^{n_i}$  for  $i = 1, 2$ , the right-hand side  $h(\cdot) \in \mathbb{R}^m$ , and the matrices  $T$  and  $W$  have orders  $n_1 \times m$  and  $n_2 \times m$ , respectively. Realizations

$\{\omega^1, \dots, \omega^S\}$ , with respective probabilities  $p^1, \dots, p^S$ , define scenarios  $\xi^s := \xi(\omega^s)$  for  $s = 1, \dots, S$ , and yield the following linear programming problem:

$$\begin{cases} \min \langle c, x_1 \rangle + \sum_{s=1}^S p^s \langle q^s, x_2^s \rangle \\ \text{s.t. } x_1 \geq 0, x_2^s \geq 0 & \text{for } s = 1, \dots, S \\ Tx_1 + Wx_2^s = h^s & \text{for } s = 1, \dots, S. \end{cases} \quad (28)$$

Decomposition by scenarios is achieved by introducing the recourse functions

$$\mathbb{Q}(x_1; \xi^s) := \begin{cases} \min \langle q^s, x_2 \rangle \\ \text{s.t. } x_2 \geq 0 \\ Wx_2 = h^s - Tx_1 \end{cases} = \begin{cases} \max \langle \pi, h^s - Tx_1 \rangle \\ \text{s.t. } W^\top \pi \leq q^s, \end{cases} \quad (29)$$

and writing (28) in the equivalent two-level formulation below:

$$\min \langle c, x_1 \rangle + \sum_{s=1}^S p^s \mathbb{Q}(x_1; \xi^s) \quad \text{s.t. } x_1 \geq 0. \quad (30)$$

The assumption of relatively complete recourse is equivalent to finiteness of the functions  $\mathbb{Q}(\cdot; \xi^s)$ , for each scenario  $s$  for all  $x_1 \geq 0$ . Complete recourse requires the condition to hold for all  $x_1$ . This stronger assumption implies that the set of dual solutions is uniformly bounded for all scenarios, somewhat simplifying some issues in the convergence analysis presented in Section 3.

The two equivalent formulations given for such recourse functions in (29) correspond to a primal and dual views (left and right, respectively). The dual view, in particular, motivated our proposal. Specifically, to “control” the multipliers, a sensible strategy would be to add a regularizing term to the objective function of the dual in (29) and, hence,

$$\begin{aligned} & \text{instead of } \mathbb{Q}(x_1; \xi^s) = \max \langle \pi, h^s - Tx_1 \rangle \quad \text{s.t. } W^\top \pi \leq q^s, \\ & \text{to consider } \mathbb{Q}^\beta(x_1; \xi^s) = \max \langle \pi, h^s - Tx_1 \rangle - \frac{\beta}{2} \|\pi\|^2 \quad \text{s.t. } W^\top \pi \leq q^s. \end{aligned}$$

For each fixed  $\beta$ , the quadratic term, making unique the solution of the dual problem (optimal multiplier of the primal), helps to prevent oscillations, and somehow stabilizes the output of the overall process with  $\beta$  variable. The relation with our initial setting (12) is explained next.

#### 4.1 The Problem to be Solved

To cast (28) as a particular instance of (12), it suffices to define the vectors

$$\begin{aligned} x &:= (x_1, x_2^1, \dots, x_2^S) \in \mathbb{R}^{n_1+n_2S}, \\ g &:= (c, p^1 q^1, \dots, p^S q^S) \in \mathbb{R}^{n_1+n_2S}, \\ b &:= (h^1, \dots, h^S) \in \mathbb{R}^{mS}, \end{aligned}$$



as well as the matrices  $H = 0 \in \mathbb{R}^{n_1+n_2S} \times \mathbb{R}^{n_1+n_2S}$ , and

$$A := \begin{bmatrix} T & W & 0 & \dots & 0 \\ T & 0 & W & \ddots & \vdots \\ T & \vdots & \ddots & \ddots & 0 \\ T & 0 & \dots & 0 & W \end{bmatrix} \in \mathbb{R}^{mS} \times \mathbb{R}^{n_1+n_2S}.$$

Given  $\beta > 0$ , it is not difficult to derive the penalized subproblems (13) for (28); specifically, we obtain:

$$\begin{cases} \min \langle c, x_1 \rangle + \sum_{s=1}^S p^s \left\{ \langle q^s, x_2^s \rangle + \frac{1}{2\beta} \|h^s - Tx_1 - Wx_2^s\|^2 \right\} \\ \text{s.t. } x_1 \geq 0, x_2^s \geq 0 \text{ for } s = 1, \dots, S. \end{cases}$$

Regarding the two-level formulation, the penalization above amounts to replacing  $\mathbb{Q}$  in (30) by the following recourse function:

$$\mathbb{Q}^\beta(x_1; \xi^s) := \min \langle q^s, x_2 \rangle + \frac{1}{2\beta} \|h^s - Tx_1 - Wx_2\|^2 \text{ s.t. } x_2 \geq 0, \quad (31)$$

whose dual formulation is

$$\mathbb{Q}^\beta(x_1; \xi^s) = \max \langle \pi, h^s - Tx_1 \rangle - \frac{\beta}{2} \|\pi\|^2 \text{ s.t. } W^\top \pi \leq q^s.$$

These primal and dual views highlight the double role of the  $\beta$ -term: a penalization in the primal becomes a regularization in the dual.

Concerning the applicability of our general convergence results to the current setting (in particular of Theorem 3.3, which establishes convergence of the dual sequence to the minimal-norm multiplier), recall that for a linear program (the case  $H = 0$ ) the key condition (18) means that the *primal* solution of problem (28) is unique. This issue was discussed in some more detail in the paragraph following (18).

## 4.2 Numerical Results

The fact that the dual second-stage problem is perturbed by a term “ $-\beta\|\pi\|^2$ ” evidently changes the dual and primal solutions, when compared to the original problem (28). Our goal is to keep close the original marginal cost, and at the same time, decrease its variance.

Theorem 3.4 describes theoretically the behavior of the mean value of regularized price signals in terms of the original optimization problem. The numerical examples below illustrate the main features of our approach.

Theoretical results in terms of variance are not simple. It is not always true that regularization reduces variance, but it happens for a large amount of problems. To make sure the approach goes in the right direction, we measured the variance and the mean value of regularized and non-regularized stochastic

problems. In addition, in (33) below we created a single index that measures the joint dynamics of variance reduction and distance to the original dual solution set, as  $\beta$  goes to zero. The index is used to make the performance profile in Subsection 4.2.2, when running the methodology on a battery of problems from the literature.

#### 4.2.1 Price Signal Analysis on an Illustrative Example

We first consider a simple instance that can be solved analytically, for checking the output. This example illustrates well our theoretical results, regarding satisfaction of condition (9), as well as convergence to the minimal-norm price (24). Take  $S = 2$  equiprobable scenarios, and let  $n_1 = n_2 = 2$ . The first-stage cost  $c \in \mathbb{R}^2$  and second-stage costs are deterministic  $q^1 = q^2 = q \in \mathbb{R}^2$ . The technology and recourse matrices in (28) are

$$T := \begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix}, \quad \text{and} \quad W := \begin{bmatrix} 2 & 0 \\ 1 & -1 \\ 1 & 2 \end{bmatrix},$$

so  $m = 3$ . The uncertain right-hand side terms are given by  $h^1 := (1, 1, 1)^\top$  and  $h^2 := (1, 0, 3)^\top$ .

Working out the algebra shows that the feasible set in (28) is completely determined by the first component of  $x_1$ , denoted by  $y \geq 0$  below. Specifically,

$$\begin{aligned} x \text{ is feasible in (28) if and only if, for some } y \geq 0, \\ x_1 := \left( y, \frac{3}{4}(1+y) \right)^\top, \quad x_2^1 := \left( \frac{1}{2}(1-y), \frac{1}{4}(1+y) \right)^\top, \\ x_2^2 := \left( \frac{1}{2}(1-y), \frac{1}{4}(5+y) \right)^\top, \end{aligned}$$

and, therefore, the following one-dimensional problem is equivalent to (28):

$$\min_{y \geq 0} \left( c_1 + \frac{3}{4}c_2 - \frac{1}{2}q_1 + \frac{1}{4}q_2 \right) y + \frac{3}{4}c_2 + \frac{1}{2}q_1 + \frac{3}{4}q_2.$$

The optimal solution of this problem is  $\bar{y} = 0$ , as long as

$$c_1 \geq -\frac{3}{4}c_2 + \frac{1}{2}q_1 - \frac{1}{4}q_2. \quad (32)$$

The primal optimum is

$$\bar{x}_1 := \left( 0, \frac{3}{4} \right)^\top, \quad \bar{x}_2^1 := \left( \frac{1}{2}, \frac{1}{4} \right)^\top, \quad \bar{x}_2^2 := \left( \frac{1}{2}, \frac{5}{4} \right)^\top.$$

To compute the optimal multiplier, recall that any normal element  $\bar{\mu} \in N_X(\bar{x})$  has all of its components null, except for the first one, because  $\langle \bar{\mu}, \bar{x} \rangle = 0$ . Therefore,

$$\bar{\mu} = -\alpha e_1 \quad \text{for some } \alpha \geq 0,$$

where  $e_j \in \mathbb{R}^6$  is the  $j$ -th canonical vector (all components are zero except the  $j$ -th, equal to 1). To check that (7) is satisfied, consider its equivalent formulation (10). Suppose  $\bar{\mu} = -\alpha e_1 \in \text{Im } A^\top$ . For condition (10) to hold, for any  $\nu \in \text{Ker } A$ , we must have that  $-\alpha \langle e_1, \nu \rangle = 0$  because the subspaces  $\text{Im } A^\top$  and  $\text{Ker } A$  are orthogonal. Since the latter (one-dimensional) subspace is generated by the vector  $s := (4, 3, -2, 1, -2, 1)^\top$ , we have that  $\langle \bar{\mu}, e^1 \rangle = -4\alpha$ , forcing  $\alpha = 0$ . It then follows that (10) and (9) hold, as claimed.

Take  $q_1 = Q = -q_2$  for some  $Q$ ,  $c_2 = 0$ , and any  $c_1 \geq Q$  (so that (32) is satisfied). Optimal Lagrange multipliers must solve the system

$$A^\top \pi = -g + \bar{\mu}, \text{ with } g = \left( c_1, 0, \frac{Q}{2}, -\frac{Q}{2}, \frac{Q}{2}, -\frac{Q}{2} \right)^\top, \text{ and } \bar{\mu} = -\alpha e_1 \text{ for } \alpha \geq 0.$$

After some algebraic manipulations, the unbounded optimal multiplier set is:

$$\mathcal{L} := \left\{ \bar{\pi} = t \frac{c_1}{2} (1, -4, 2, -3, 4, 2)^\top \mid t \geq 1 \right\}.$$

Hence,  $t = 1$  gives the minimal-norm element for which  $\mathbb{E}[\hat{\pi}] = \frac{c_1}{2} (-1, 0, 2)^\top$ .




Fig. 1: Unbounded set of optimal multipliers in mean (the line), the element with minimal norm (the plus sign), the mean multiplier found for  $\beta = 0$  (the cross), the mean multiplier estimates for different values of  $\beta > 0$  (the triangles), and the origin (the dot)

Applying our approach with several decreasing values of  $\beta$  gives the multiplier estimates  $\pi^\beta \in \mathbb{R}^6$  with the mean  $\mathbb{E}[\pi^\beta] \in \mathbb{R}^3$ . The line in Figure 1 shows a portion of the mean optimal multiplier set,  $\left\{ t \frac{c_1}{2} (-1, 0, 2)^\top \mid t \geq 1 \right\}$ . The dot represents the origin in  $\mathbb{R}^3$ , the plus sign  $\mathbb{E}[\hat{\pi}]$ , the minimal-norm multiplier in mean value, to which the mean values  $\mathbb{E}[\pi^\beta]$ , represented with triangles, converge as  $\beta \rightarrow 0$ . Finally, the cross displays  $\mathbb{E}[\bar{\pi}]$ , the mean multiplier found when solving (28), whose norm is larger than the minimal one.

#### 4.2.2 Combined Index of Variance and Mean Value

Performance profiles [19] are useful tools to benchmark different methods on a fair basis. For a battery of two-stage stochastic linear programming problems, we compare the expected value of the multipliers, obtained as follows:

- Solving (28) in its two-level formulation; (30) with recourse function (29), by a proximal bundle method [20]; see also [21, Ch. 10.3].
- Our proposal, i.e., solving for decreasing values of  $\beta$  several instances of (30), with recourse function (31).

All the tests were ran in Matlab R2016, on an Intel Core i5 computer with 2.4 GHz, 4 cores and 4 GB RAM, running under Ubuntu 18.04.1 LTS and using Gurobi 5.6 optimization toolbox for Matlab.

The battery comprises 50 problems, for which 10 independent instances, each one with 50 scenarios, were created. The considered two-stage stochastic problems are of the form (28) with uncertainty only on the right-hand side  $h \in \mathbb{R}^m$ , independently and normally distributed. The expectation and standard deviation of the considered distribution is problem-dependent, and proportional to the vector  $\frac{c}{2}$ . The problem dimension ranges are  $n_1 \in \{20, 40, 60\}$ ,  $n_2 \in \{30, 60, 90\}$ , and  $m \in \{20, 40, 60\}$ . For more details we refer to [22, 10].

The test has a total of 500 runs, labeled  $P = 1, \dots, 500$ . With the purpose of doing a performance profile, we compute

$$\|\mathbb{E}[\pi_P^{\text{best}}]\| := \arg \min \left\{ \|\mathbb{E}[\pi_P^\beta]\| : \beta \in \{0, 0.1, 0.2, \dots, 0.5\} \right\},$$

and define, for problem  $p$  and parameter  $\beta \geq 0$ , the following index:

$$c_P^\beta := \frac{\|\text{Var}[\mathbb{E}[\pi_P^\beta]]\|}{\|\text{Var}[\pi_P^{\text{best}}]\|} + \left( 1 - \frac{\|\mathbb{E}[\pi_P^\beta]\|}{\|\mathbb{E}[\pi_P^{\text{best}}]\|} \right). \quad (33)$$

Here,  $\pi_P^0$  corresponds to  $\bar{\pi}$  in our previous notation, while  $\pi_P^\beta$  is the Lagrange multiplier computed for problem  $P$  with regularization parameter  $\beta \geq 0$ . The corresponding performance profile is given in Figure 2.




Fig. 2: Combined gains in expected value and variance for the dual variable.

In terms of the combined index, and as expected, the multiplier of the original problem ( $\beta = 0$ ) performs worse, confirming the empirical observation that in general  $Var[\pi_P^\beta] \leq Var[\pi_P^0]$ . For this set of runs, the value  $\beta = 0.1$  (dashed line with circles) seems to give a good compromise between stability of the mean multiplier, and approximation of the minimal-norm multiplier.

For completeness, we present in Figure 3 a profile for the first-stage variable, measuring its performance with the index

$$\tilde{c}_P^\beta := \left( 1 - \frac{\|\mathbb{E}[x_{1P}^\beta]\|}{\|\mathbb{E}[x_{1P}^0]\|} \right),$$

defined for  $\beta > 0$  (comparing variances is not sound in our approach, as there is no “stabilization” of the primal variables). It can be seen in the graph that the best value for  $\beta$  in the dual performance profile in Figure 2, that is  $\beta = 0.1$  (dashed line with circles), behaves reasonably well in the primal variable.




Fig. 3: Progression of expected value of first-stage variable

## Conclusions

In many applications, dual variables are an important output of the decision/solution process, due to their role as price signals. When dual solutions are not unique, different solvers or different computers, even different runs in the same computer if the problem is stochastic, end up with different price indicators. Even though all of such values are mathematically correct, the fact that the obtained dual variable varies among many possibilities makes unreliable any economic analysis based on marginal prices. We have presented an approach that yields reliable indicators, by providing the minimal-norm multiplier. Our proof-of-concept computational experience shows the benefits of the methodology for two-stage stochastic linear programs.

The best choice for the penalization/regularization parameter  $\beta$  is clearly problem dependent. Somewhat similarly to the solution concept called *compromise decision* in [23], but adopting a dual point-of-view, the performance index proposed in (33) aims at measuring bias and variance, in multiple replications of sampling-based approximations of two-stage stochastic programs.

We observe empirically that our approach yields a significant reduction in the variance of the dual solutions (optimal Lagrange multipliers). A topic of ongoing research is to develop a quantitative stability analysis, along the lines of [24], but on the dual variables; see also [25,26].

## Acknowledgements

This research was supported by a CIFRE contract between ENGIE and Université de Paris Sorbonne, France. The first author is joint PhD candidate of IMPA – Instituto de Matemática Pura e Aplicada (Brazil) and Université de Paris Sorbonne (France). The second author’s research is also partially supported by CNPq Grant 303905/2015-8 and by FAPERJ. Research of the third author is also supported in part by CNPq Grant 303724/2015-3, by FAPERJ Grant 203.052/2016, and by the Russian Foundation for Basic Research Grant 19-51-12003 NNIOa.

The authors are grateful to W. de Oliveira for providing the Matlab code for the test functions used in the performance profiles.

## References

1. Shapiro, A., Dentcheva, D., Ruszczyński, A.: Lectures on Stochastic Programming: Modeling and Theory. Society for Industrial and Applied Mathematics, Philadelphia (2009)
2. Kleywegt, A., Shapiro, A., Homem de Mello, T.: The sample average approximation method for stochastic discrete optimization. *SIAM J. Optim.* 12, 479-502 (2001/02)
3. Eisner, M. J., Olsen, P.: Duality for stochastic programming interpreted as L.P. in  $L_p$ -space. *SIAM J. Appl. Math.* 28, 779-793 (1975)
4. Rockafellar, R. T., Wets, R.: Stochastic convex programming: Basic duality. *Pacific J. Math.* 62, 173-195 (1976)
5. Rockafellar, R. T., Wets, R.: Stochastic convex programming: singular multipliers and extended duality singular multipliers and duality. *Pacific J. Math.* 62, 507-522 (1976)
6. Slyke, R.M., Wets, R.: L-shaped linear programs with applications to optimal control and stochastic programming. *SIAM J. Appl. Math.* 17, 638-663 (1969)
7. Birge, J., Louveaux, F.: A multicut algorithm for two-stage stochastic linear programs. *Euro. J. Oper. Res.* 34, 384-392 (1988)
8. Ruszczyński, A.: Some advances in decomposition methods for stochastic linear programming. *Ann. Oper. Res.* 85, 153-172 (1999)
9. Zakeri, G., Philpott, A., Ryan, D.: Inexact cuts in Benders decomposition. *SIAM J. Optim.* 10, 643-657 (2000)
10. Oliveira, W., Sagastizábal, C., Scheimberg, S.: Inexact bundle methods for two-stage Stochastic Programming. *J. Optim.* 21, 517-544 (2011)
11. Fábíán, C., Wolf, C., Koberstein, A., Suhl, L.: Risk-averse optimization in two-stage stochastic models: Computational aspects and a study. *SIAM J. Optim.* 25, 28-52 (2015)
12. Ackooij, W., Malick, J.: Decomposition algorithm for large-scale two-stage unit-commitment. *Ann. Oper. Res.* 238, 587-613 (2016)
13. Rockafellar, R., Wets, R.: Variational Analysis. Springer Verlag, Berlin (1998)
14. Solodov, M.: Constraint qualifications. In *Wiley Encyclopedia of Operations Research and Management Science*, James J. Cochran, et al. (editors), John Wiley & Sons, Inc., 2010.
15. Izmailov, A., Solodov, M.: Newton-type Methods for Optimization and Variational Problems. Springer, New York (2014)
16. Nocedal, J., Wright, S.: Numerical Optimization. Springer, New York (2006)
17. Fiacco, A.V., McCormick, G.: Nonlinear Programming: Sequential Unconstrained Minimization Techniques. John Wiley & Sons, New York (1968).

18. Frank, M., Wolfe, P.: An algorithm for quadratic programming. *Naval research logistics quarterly*. *Nav. Res. Logis. Quart.* 3, 95-110 (1956)
19. Dolan, E.D., Moré J.J.: Benchmarking optimization software with performance profiles. *Math. Program.* 91, 201-213 (2002)
20. Lemaréchal, C., Sagastizábal, C.: Variable metric bundle methods: from conceptual to implementable forms. *Math. Program.* 76, 393-410 (1997)
21. Bonnans, J.F., Gilbert, J.C., Lemaréchal, C., Sagastizábal, C.: *Numerical Optimization. Theoretical and Practical Aspects*. Springer-Verlag, Berlin (2006)
22. Deak, I.: Two-stage stochastic problems with correlated normal variables. *Computational experiences*. *Ann. Oper. Res.* 142, 79-97 (2006)
23. Sen, S., Liu, Y.: Mitigating uncertainty via compromise decisions in two-stage stochastic linear programming, Variance reduction. *Oper. Res.* 64, 1422-1437 (2016)
24. Liu, Y., Römisch, W., Xu, H.: Quantitative stability analysis of stochastic generalized equations. *SIAM J. Optim.* 24, 467-497 (2014)
25. Dentcheva, D., Römisch, W.: Differential stability of two-stage stochastic programs. *SIAM J. Optim.* 11, 87-112 (2000)
26. Römisch, W.: Stability of stochastic programming problems. *Handb. Oper. Res. Man. Sci.* 10, 483-554 (2003)