



HAL
open science

Influenza Mortality in French Regions after the Hong Kong Flu Pandemic

Florian Bonnet, Hippolyte d'Albis, Josselin Thuilliez

► **To cite this version:**

Florian Bonnet, Hippolyte d'Albis, Josselin Thuilliez. Influenza Mortality in French Regions after the Hong Kong Flu Pandemic. 2022. halshs-03763371

HAL Id: halshs-03763371

<https://shs.hal.science/halshs-03763371>

Preprint submitted on 29 Aug 2022

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



PARIS SCHOOL OF ECONOMICS
ECOLE D'ECONOMIE DE PARIS

WORKING PAPER N° 2022 – 27

Influenza Mortality in French Regions after the Hong Kong Flu Pandemic

Florian Bonnet
Hippolyte d'Albis
Josselin Thuilliez

JEL Codes:
Keywords:

anr®
agence nationale
de la recherche
AU SERVICE DE LA SCIENCE



Influenza Mortality in French Regions after the Hong Kong Flu Pandemic

Florian Bonnet¹ – Hippolyte d'Albis² – Josselin Thuilliez³

Influenza mortality has dramatically decreased in France since the 1950s. Annual death rates peaked during two pandemics: the Asian flu (1956-57) and the Hong-Kong flu (1969-1970). This study's objective is to evaluate whether the second pandemic created a structural change in the dynamics of influenza mortality in France. We employ a new database on influenza mortality since 1950 at the subnational level (90 geographic areas) to estimate statistical models to find whether a structural change happened and to explain the differences in mortality rates across geographic areas. Influenza mortality increased between 1950 and 1969, and decreased from 1970 onward. The Hong-Kong flu is identified as the event of a structural break. After the break, geographical differences are less explained by regional characteristics such as income, density or aging ratio. Hong Kong flu was found to be associated with a major change in influenza mortality in France. Change in health practices and policies induced a decline in mortality that started in 1970, just after the pandemics. The health benefits are notably important for senior citizens and for the poorest regions.

¹ INED (Institut National d'Etudes Démographiques). Orcid: 0000-0001-6125-3317.

² Paris School of Economics, CNRS (Centre National de la Recherche Scientifique). Orcid: 0000-0002-6409-4320.

³ Centre d'économie de la Sorbonne, CNRS (Centre National de la Recherche Scientifique). Orcid: 0000-0001-5392-0917.

1. Introduction


The Covid-19 pandemic has strongly afflicted France. Yet, little attention has been given to the health outcomes following previous pandemics. Pandemics are dramatic shocks which have the ability to positively transform health systems and practices over the long run, as they generate political and economic support for medical innovation and healthcare improvement (Dehner 2010; Morens, Taubenberger and Fauci 2009; Jordà, Singh and Taylor 2020; Iskander et al. 2013). Before Covid-19, and aside from the global HIV epidemic, only two pandemics severely affected France after WWII: the Asian flu (1956-57) and the Hong-Kong flu (1969-1970). Death tolls of these pandemics were much lower than that of the current crisis, even adjusted for population and age distribution changes: 33,232 deaths from the Asian flu and 32,062 deaths from the Hong-Kong flu. The main difference is that in earlier pandemics, the mortality rates of infants and young adults were higher than that of today.

In the aftermath of these two pandemics, international health authorities and industrialized countries' governments carefully crafted responses, shared information, established surveillance, strengthened vaccine studies, and implemented vaccination campaigns against influenza (Tyrrell et al. 1970; Carrat and Valleron 1995; Sands, Mundaca-Shah and Dzau 2016; Viboud et al. 2005; Aranzazu 2013; Vagneron 2015). The aim of our study is to evaluate whether this was effective and, more precisely, whether these pandemics created a structural break in the trend line of France's flu mortality. To do so, we examine a unique historical database that tracks flu mortality by region. Our contribution is threefold. First, we provide a refined description of influenza pandemics and seasonal influenza in 90 French geographical units from 1950 to 2015. Second, we establish that winter 1969-1970 does indeed represent a structural break in the spread and control of seasonal and pandemic influenza in France, especially for the elderly, confirming a major turning point in France's flu management. Third, we show that, after 1970, influenza mortality became regionally random, no longer dependent on income, density, and the share of elderly people.

The Hong Kong flu led to a sharp increase in the death rate of flu: in 1969, it was 30 per 100,000 in France, compared to an average of 14 per 100,000 between 1965 and 1974. Moreover, **Figure 1** reveals

that geographic variations were significant: death rates were lower in the North of France (with a minimum of 12.5 per 100,000 in Seine) and higher in the South-West (with a maximum of 76.4 per 100,000 in Creuse). Despite a multilevel response to the Hong Kong flu aimed at preventing future such health shocks, the current crisis reveals that France was not fully prepared. However, we show that pandemics can change health practices for the better, as the Hong-Kong flu episode is found to be associated with a reduction in both mortality and inequalities therein across regions.

Figure 1. Map of flu death rates by departments in 1969 (deaths per 100,000 population).



2. Methods

2.1. Data

We gathered original death data by cause, geographical unit (*département* in French, hereby “department”), and sex for the period 1950-2015. The data was retrieved from the archives of the *Statistique des Causes de Décès* published by *Institut National des Statistiques et des Etudes Economiques* (INSEE) for the period 1950-1978, and from *Institut National de la Santé et de la Recherche Médicale* (INSERM) for the period 1979-2015. Figures come from the International Classification of Diseases short list. It follows the classification of the 6th revision from 1950 to 1957, the 7th revision from 1958 to 1967, the 8th revision from 1968 to 1978, the 9th revision from 1979 to 1999 and the 10th revision from 2000 to 2015. We retrieved deaths due to influenza as well as total deaths by department and sex for the entire period.

Our study focuses on mainland France, and does not include overseas territories. We use the classification of 90 departments that prevailed before 1968, but we also provide analysis by aggregating the data into 22 geographical units known as regions (hereby “regions”).

For the statistical analysis, we also use population data by department, sex and age at 1st January. These populations were calculated from age-specific populations in census years and annual vital statistics (age-specific deaths and births), using a methodological protocol close to that of the Human Mortality Database (Bonnet 2020; Barbieri et al. 2015). To obtain the populations on January 1 of each year of an intercensal period, the author adjusts the populations at the initial census by both observed deaths and by an estimate of the migration flow since the initial census. The overall intercensal migration flow is estimated by comparing the observed population and the theoretical population at the final census. The theoretical population is equal to the initial census population minus the observed deaths during the intercensal period. The intercensal migration flow is then distributed temporally in proportion to the time elapsed since the initial census. To compute death rates in year t , we use mean of population at 1st January of year t and $t+1$.

We also use income data by department (Bonnet and Sotura 2021; Bonnet, d’Albis, and Sotura 2021). These incomes were calculated using tax tabulations for the period 1960-1969 and 1986-2015 and aggregate income tax statistics for the periods 1922-1959 and 1970-1985. This income includes public transfers related to the pension system, but does not take into account taxes due or public subsidies provided to households.

2.2. Statistical Analysis

We aim to identify whether and when there was a structural break in the control of influenza epidemics in France. We use a Poisson model, that prevents from negative values for death rates, and regress deaths on a continuous year variable, a post-year dummy, and the interaction of post x year dummy, in addition to department-specific time trend, department fixed-effects, and specific controls (population density per km², per capita income in 2015 euros, and the share of people aged 65 and more in the population). To take into account the difference in department sizes, we use population as an offset in this regression. We repeat the procedure, allowing the post-year dummy to vary from 1960 to 1980. The year with maximum in log-likelihood is considered to be the year of a structural break in the data. This test is similar to the Quandt likelihood ratio test that has been shown to be a reliable test for structural breaks in the case of unknown break points (Quandt, 1960; Andrews, 1993; Piehl et al., 2003).

We also perform a spatial analysis. We examine the differences in deaths across department using Poisson model and three specific variables: population density per km² (D), income per capita in 2015 euros (Inc) and the share of people aged 65 and more in the population (Sh_d^{65+}). For each year of the period 1950-2015, we estimate the coefficients of the following equation:

$$\log(m_d^{Flu}) = \alpha + \beta_1 D_d + \beta_2 I_d + \beta_3 Sh_d^{65+} + \varepsilon_d, (1)$$

where d is the department and ε_d is the difference between observed and estimated mortality due to flu. If positive, it is defined as excess mortality. To compute confidence intervals around our estimates, we consider over-dispersion in the model. Finally, we compute pseudo R² of our model for each year as the square of the correlation between estimated and observed departmental values. All our computations are

done using R. software. **Table 1** presents descriptive statistics for variables used in Equation (1) and specific years.

Table 1. Descriptive statistics.

	1950	1960	1970	1980	1990	2000	2010
Flu death rate	5.25 (2.98)	32.03 (11.96)	18.1 (8.67)	2.41 (1.67)	5.69 (4.84)	3.45 (2.17)	0.24 (0.27)
Density	183.21 (1069.73)	205.11 (1209.71)	233.26 (1394.3)	234.86 (1342.6)	239.67 (1334.17)	245.53 (1349.34)	263.25 (1452.56)
Income per adult	2407.63 (615.77)	3855.42 (788.22)	6720.35 (1153.78)	9896.5 (1328.44)	11327 (1415.24)	13402.45 (1487.54)	15488.09 (1465.79)
Share of 65+	12.46 (2.07)	12.56 (2.09)	13.94 (2.53)	15.6 (3.15)	15.66 (3.2)	18.07 (3.44)	19.03 (3.24)

Notes: We provide mean and standard deviation (in parenthesis) for 90 departmental values. Flu death rates are per 100,000, income per capita are in 2015 euros.


3. Results

The test for a trend break in mortality between 1950 and 2015 identifies 1968 as the year with the maximum in log-likelihood. Values are roughly similar for each year of the period 1968-1971, and highly decrease in 1972. There are other spikes related to influenza outbreaks (1974 and 1977), but they are smaller in magnitude. We note that sensitivity testing of our regression does not affect our main conclusions. We retain 1970 as the year of the structural break for our subsequent analyses since it is the end of the Hong-Kong flu and the central year of the period 1968-1971.

We present a range of graphs to illustrate our empirical results. **Figure 2** shows flu mortality by region from 1950 to 2015. The red lines provide estimates of mortality trends for the periods 1950-1970 and 1970-2015 using Poisson model. The Hong Kong flu period (i.e. 1968-1970) appears in grey. The last figure provides flu specific mortality at the national level while **Figures A1, A2 and A3** in Appendix provides the flu mortality by department. In France, we find a sharp decline in flu mortality between 1950-1960 and 2005-2015, of a magnitude between 84% and 98%. This decrease is sustained over time and converges towards 5 per 100,000 on average in 2015. Importantly, we observe that almost no region

or department exhibits, before 1970, a negatively sloped mortality trend. Mortality was, in fact, increasing before the Hong Kong flu and decreased thereafter.


Figure 2. Flu death rates by region (deaths per 100,000 population).



Notes: This figure plots flu mortality rates by year and region for the period 1950-2015. The red lines provide estimates of mortality trends for the periods 1950-1970 and 1970-2015 using Poisson model. Hong Kong flu (1968-1970) is shown in grey.

Figure 3 plots mean flu death rates in France’s 90 departments between 1950 and 1968 against the variation in mean flu death rates between 1950-1968 and 1970-2015. The slope of the relationship is -0.64. This reveals a strong convergence across geographical units; in addition to across-the-board declines in flu-specific mortality, the initially-high-mortality departments essentially caught up to the initially-low-mortality departments. This convergence is associated with a reduction in geographical inequalities, as documented for the general mortality (Bonnet and d’Albis 2020).


Figure 3. Declines in flu specific mortality across departments (deaths per 100,000 population).



Notes: This figure plots mean death rates of flu between 1950 and 1968, and the variation in mean death rates of flu between 1950-1968 and 1970-2015 for the France’s 90 departments. The results of the regression is plotted in red. The value of the slope is equal to -0.65.

Figure 4 provides a decomposition by age group of flu specific mortality in France. These results suggest that the break caused by the Hong Kong flu affected only adults, particularly senior citizens. The decline in flu mortality for those under age 25 started much earlier and 1970 death rates for those ages 1 to 44 were already very low. Thus, the shock of the Hong Kong flu appears to have induced the healthcare system to devote more attention to flu mortality of the elderly. The death rate of those 65+ shifted from an average of 200 deaths per 100,000 in the 1960’s to an average of 30 per 100,000 in the 1980’s. The response to the Covid-19 crisis echoes such consideration, initiating a historic worldwide effort to predominantly protect the elderly. We also notice that the structural break holds for both men and women, the death rate of the latter being slightly higher due to differences in age structure.


Figure 4. Death rates of flu by age groups and sex in France (deaths per 100,000 population).



Notes: This figure plots death rates of flu by year, age group and sex for the period 1950-2015. The red lines provide estimates of mortality trends for the periods 1950-1970 and 1970-2015 using a Poisson model. The period of the Hong Kong flu (i.e. 1968-1970) appears in grey.

Figure 5 plots the coefficients estimated in the linear regression characterizing geographical death rates over time. Notably, it reveals that per capita income explained flu mortality rates in some major flu outbreaks prior to and during Hong Kong flu: controlling for population density and the share of people aged 65+, those with higher incomes had lower mortality rates. This disadvantage for the poorest departments was reduced after the Hong Kong flu. Thus, improved health practices benefited the poorest departments, which had suffered more than others from the flu. Overall, we note that, after the Hong Kong flu, differences in these characteristics explain decreasing amounts of variation in geographical differences in flu mortality: the pseudo R^2 associated with each annual regression decreases to close to 0 in 2010 after reaching its maximum (0.45) in 1969. We interpret this as flu mortality becoming more regionally stochastic.

Figure 5. Explanations of differences in department flu death rates.



Notes: This figure plots the annual coefficients of equation (1), and the 95% confidence intervals of these values. This figure represents the pseudo R2s of the annual regressions too. Values statistically different from 0 are in red. The period of the Hong Kong flu (i.e. 1968-1970) appears in grey.

4. Discussion

The Hong Kong flu pandemic started in July 1968 and lasted until 1970. Despite the staggering death toll, the Hong Kong flu pandemic and its repercussions are surprisingly poorly documented in France (Flahault and Zylberman 2010). Beyond natural immunity which may have played a role that is difficult to assess in the medium run (Jackson, Vynnycky and Mangtani 2010), what could be behind such a strong decline of influenza across all departments in France after 1970? It is difficult to identify any major progress in hygiene at that time (Flahault and Zylberman 2010). The discovery of single-use handkerchiefs dates back to the 1930s and massive use of hydro-alcoholic gel, for instance, dates much later. Surveillance has indeed been improved, but this progress seems to have taken place mainly during

the 1980s. Finally, there is also little chance that France's influenza is strongly correlated with economic growth or crises (Ruhm 2000; Brüning and Thuilliez 2019).

As we examine what caused the decline, the distinction between pandemic and seasonal flu is important (Brüning and Thuilliez 2019). France has not experienced a severe pandemic from 1969 onwards, either because the world was better prepared, or just by chance. Put differently, reduced seasonal flu mortality after 1970 may not have been influenced by pandemics, but rather part of a pattern of seasonal flu (Carrat and Valleron 1995; Rizzo et al. 2007; Guérin 2007; Gottfredsson et al. 2008; Valleron et al. 2010; Hannoun 2013; Brès 1980). **Figure A4 and A5** in Appendix hence test for the sensitivity of our results to the presence of pandemics and provide the change in the slope when we exclude the 1957 and 1968 pandemics, assuming a value for these particular years as equivalent to the average mortality before 1970. The difference between the slopes gives us the contribution of pandemics to the average mortality trend. When both pandemics are eliminated, the slopes before 1970 (in blue) move downward for all regions and age groups and mostly for those age groups mostly affected by pandemics, but the structural break remains apparent. This confirms the influence of pandemics in the change in flu mortality.

A change in vaccination practices could be amongst the most plausible explanations of the structural break (Meslé 2010). Lacking historical data, we were unable to find reliable figures on the production of influenza vaccines in France from 1950 to 1976. However, the WHO delivered vaccination recommendations in the early 1970s and various public health organizations were also founded at that time (Brès 1980). **Figure A6** in Appendix provides tentative estimates of influenza vaccination in France from 1950 to 2015 from unpublished data and they indicate an increase after 1970. We do not have data on vaccination at sub-national level, by age groups or SES over this period of time, to the best of our knowledge. Vaccination rate before 1977 is therefore estimated according to 4 scenarios because we do not have data for this period in France. However, influenza vaccination principles were already well known at that time and the population was aware of vaccination benefits since the Pasteurian revolution. **Figure A7** in Appendix illustrates that the epidemic was well covered by the media (MacCarthy 1969), as exemplified in an interview of Robert Boulin, the Health Minister, who got vaccinated and strongly

recommend it to the population. Another article documents the high demand pharmacies faced for the vaccines. In addition, **Figure A8** in Appendix shows that flu-specific mortality for senior citizens is highly correlated with vaccination in the recent period, as we observe a decline in vaccination, notably during the 2009 A(H1N1) influenza pandemic (Vaux et al. 2011; Blondel et al. 2012), and an increase in mortality since 2010. Other technological and medical advances, some related to co-morbidities or complications of influenza, may also have played a role. Pneumococcal vaccines, for example, help to reduce influenza mortality (Fedson 1998). Nevertheless, in France, whether the public is willing to accept vaccination when it is available remains important (Schwarzinger et al. 2010; Peretti-Watel, Seror et al. 2020).

Studies on this topic may also include pneumonia deaths. The counterpart of **Figure 2** for pneumonia related death is provided in **Figures A9a and A9b** in Appendix. The patterns are quite different for that of influenza. Interestingly, we observe a hump in the 90's, certainly due to the HIV-AIDS epidemics. The decline also coincides with the introduction of highly active antiretroviral therapies (McCullers 2013; Metersky et al. 2012; Shrestha et al. 2015). Post-influenza bacterial pneumonia is a major cause of morbidity and mortality associated with both seasonal and pandemic influenza virus illnesses. However, understanding the role that influenza plays on the population-level epidemiology of bacterial pneumonia and, reciprocally, the role of bacterial pneumonia complicating influenza, remains a challenge, the reason why we prefer to keep both causes of deaths separated (Feikin et al. 2004; Franquet and Domingo 2022; Roux et al. 2014). We also replicate **Figure 2** for all-cause mortality in **Figure A10** in Appendix. There is no break in this period, which reinforces our argument about a specificity of the flu.

A “dry tender” effect may mitigate our story on structural changes in the short-run (Rizzi, Søggaard and Vaupel 2021). Indeed, harvesting could partly explain the fall in mortality, and also the decline in the role of income after 1970. The “Dry Tinder” effect has been well documented by epidemiologists to describe a situation where a soft year creates a larger group of susceptible individuals the following year. We recognize that such a selection effect could partly explain the fall in mortality in the short-run. However, after a few years, the most vulnerable age groups (infants and elderly) are replaced by new

cohorts and the effect is less likely to explain the change we observe. Moreover, **Figure A6 in Appendix** shows that vaccine uptake increases substantially after 1969.

We also use our model over the entire period (rather than year by year) to predict the values at the national level. The difference between the observed values and the predicted values is excess deaths. We reported this in **Figure A11 in Appendix**. We can see clearly two periods, before and after the Hong-Kong flu. Years of excess deaths are associated to the main epidemics, and from the 1970s onwards, we observe a permanent under-mortality. 2009 H1N1 pandemic had relatively low mortality effects in France. However, it is true that the 2009 H1N1 pandemic changed public attitudes towards vaccination in France (Lemaitre and Carrat 2010; Lemaitre et al. 2012; Boëlle et al. 2020; Fuhrman et al. 2010). We note that in mainland France, the 2010-2011 season was characterized in the community by an influenza epidemic of moderate intensity, which occurred between late December 2010 and mid-February 2011. Severe cases admitted to the ICU had epidemiological characteristics and lethality comparable to those observed during the 2009-2010 pandemic and vaccine hesitancy increased in France after the H1N1 pandemic (Schwarzinger et al., 2010; Ward et al. 2019; Blanchon 2011).

Lastly, this paper has some limitations. First, Figure 2 shows subnational heterogeneity of levels and trends. When pooling all regions, we obtain a break in 1968, and when we separate our dataset by region we obtain 1968 or 1969 as a break in 18 regions out of 22 (2 for 1970, 1 for 1967, 1 for 1974). Therefore, overall our results seem to be consistent across regions despite different magnitudes in the breaks observed at the regional level. They mostly confirm that the break is at least in 1968 and not before this date except for one region. Disaggregating even more at the department level may of course lead to more heterogeneity. Second, we use crude death rates in this paper. Crude death rates describe how many people die out of a given population over a specific time period but take no account of the population's age distribution when age-standardized measurements of mortality are not available. We use as a control in each model the share of the population aged 65 and over, ages for which influenza mortality is highest. We could control for the share of each five-year age group. Because we stratify the data and present the results without aggregating the stratum-specific information over the strata, we preferred to use crude death rates. Third, our graphical results and regression methods are not fully

causal but our approach and our robustness tests concerning the date of the break are suggestive of a causal explanation.

In conclusion, we found the Hong Kong flu to be associated with a major change in influenza mortality in France. Using a unique dataset of 90 geographical units since 1950, we found a decline in mortality beginning in 1970, just after the major pandemics. The change was most notable for senior citizens and the poorest regions. These facts are useful to put the recent Covid-19 crisis in perspective. First, they reveal that health behaviors and policies are impacted by these dramatic events (Morse 2007); we can thus expect profound changes to our health system in the coming years. Second, they suggest that the value of elderly life became more important in the late 1960's, a paradox for a period that is usually perceived as a golden age for youth. Finally, the flu's geographical differences were strongly reduced after the 1970's, with mortality rates that became uncorrelated to regional mean income. Geographical prevalence became a random event, and pandemics may thus hit regions differently, as we have seen in the Covid-19 crisis. The Covid-19 pandemic is thus a good example that seems to indicate that preparedness and former pandemics experience played a key role in the crisis management. However, better preparedness cannot hinder the probability that a pandemic occurs, all the more if the disease was not circulating before its sudden emergence.

References

- Andrews, D. (1993). Tests for parameter instability and structural change with unknown change point. *Econometrica*, 61: 821-856.
- Aranzazu, A. (2013). Le réseau mondial de surveillance de la grippe de l'OMS. Modalités de circulation des souches virales, des savoirs et des techniques, 1947-2007. *Sciences sociales et sante*, 31: 41-64.
- Barbieri, M., Wilmoth, J. R., Shkolnikov, V. M., Gleijeses, D., Jasilionis, D., Jdanov, D., Boe, C., Riffe, T., Grigoriev, P. and Winant, C. (2015). Data resource profile: the human mortality database (HMD). *International journal of epidemiology*, 44: 1549-1556.
- Bellia, C., Setbon, M., Zylberman, P., and Flahault, A. (2013). Healthcare worker compliance with seasonal and pandemic influenza vaccination. *Influenza and Other Respiratory Viruses*, 7: 97-104.
- Blanchon, T. (2011). Surveillance épidémiologique et virologique de la grippe en France, saison 2010-2011.
- Blondel, B., Mahjoub, N., Drewniak, N., Launay, O., and Goffinet, F. (2012). Failure of the vaccination campaign against A(H1N1) influenza in pregnant women in France: Results from a national survey. *Vaccine*, 30: 5661-5665.
- Boëlle, P.-Y., Souty, C., Launay, T., Guerrisi, C., Turbelin, C., Behillil, S., Enouf, V., Poletto, C., Lina, B., van der Werf, S., Lévy-Bruhl, D., Colizza, V., Hanslik, T. and Blanchon, T. (2020). Excess cases of influenza-like illnesses synchronous with coronavirus disease (COVID-19) epidemic, France, March 2020. *Eurosurveillance*, 25:2000326.
- Bonnet, F. (2020). Computations of French lifetables by department, 1901--2014. *Demographic Research*, 42: 741-762.
- Bonnet, F., and d'Albis, H. (2020). Spatial Inequality in Mortality in France over the Past Two Centuries. *Population and Development Review*, 46: 145-168.
- Bonnet, F., and Sotura, A. (2021). Regional Income Distributions in France, 1960-2018. *Banque de France Working Papers*, 832.
- Bonnet, F., d'Albis, H., and Sotura, A. (2021). Les inégalités de revenu entre les départements français depuis cent ans/Income Inequality across French Departments over the Last 100 Years. *Economie et Statistique*, 526: 49-70.
- Brès, P. (1980). Surveillance épidémiologique des gripes humaines dans le monde: Le programme de l'OMS. *Comparative Immunology, Microbiology and Infectious Diseases*, 3: 33-43.
- Brüning, M., and Thuilliez, J. (2019). Mortality and Macroeconomic Conditions: What Can We Learn From France? *Demography*, 56: 1747-1764.
- Carrat, F., and Valleron, A. J. (1995). Influenza mortality among the elderly in France, 1980-90: how many deaths may have been avoided through vaccination? *Journal of Epidemiology & Community Health*, 49: 419-425.


- Dehner, G. (2010). WHO Knows Best? National and International Responses to Pandemic Threats and the "Lessons" of 1976. *Journal of the History of Medicine and Allied Sciences*, 65: 478-513.
- Fedson, D. S. (1998). Pneumococcal Vaccination in the United States and 20 Other Developed Countries, 1981–1996. *Clinical Infectious Diseases*, 26: 1117-1123.
- Feikin, D. R., Feldman, C., Schuchat, A., and Janoff, E. N. (2004). Global strategies to prevent bacterial pneumonia in adults with HIV disease. *The Lancet Infectious Diseases*, 4: 445-455.
- Flahault, A., & Zylberman, P. (2010). Influenza pandemics: past, present and future challenges. *Public Health Reviews*, 32: 319-340.
- Franquet, T., & Domingo, P. (2022). Pulmonary Infections in People Living with HIV. *Radiologic Clinics of North America*, 60: 507-520.
- Fuhrman, C., Bonmarin, I., Paty, A. C., Duport, N., Chiron, E., Lucas, E. and others. (2010). Severe hospitalised 2009 pandemic influenza A (H1N1) cases in France, 1 July-15 November 2009. *Eurosurveillance*, 15: 19463.
- Gottfredsson, M., Halldórsson, B. V., Jónsson, S., Kristjánsson, M., Kristjánsson, K., Kristinsson, and others. (2008). Lessons from the past: Familial aggregation analysis of fatal pandemic influenza (Spanish flu) in Iceland in 1918. *Proceedings of the National Academy of Sciences*, 105: 1303-1308.
- Grais, R. F., Hugh Ellis, J., and Glass, G. E. (2003). Assessing the impact of airline travel on the geographic spread of pandemic influenza. *European Journal of Epidemiology*, 18: 1065-1072.
- Guérin, N. (2007). Histoire de la vaccination: de l'empirisme aux vaccins recombinants. *La Revue de Médecine Interne*, 28: 3-8.
- Hannoun, C. (1985). Les Vaccins contre la grippe. *Annales de l'Institut Pasteur / Virologie*, 136: 553-559.
- Hannoun, C. (1997). Le vaccin antigrippal d'aujourd'hui et de demain. *Virologie*, 1: 121-31.
- Hannoun, C. (2013). The evolving history of influenza viruses and influenza vaccines. *Expert Review of Vaccines*, 12: 1085-1094.
- Iskander, J., Strikas, R. A., Gensheimer, K. F., Cox, N. J., and Redd, S. C. (2013). Pandemic Influenza Planning, United States, 1978–2008. *Emerging Infectious Diseases*, 19: 879-885.
- Jackson, C., Vynnycky, E., and Mangtani, P. (2010). Estimates of the Transmissibility of the 1968 (Hong Kong) Influenza Pandemic: Evidence of Increased Transmissibility Between Successive Waves. *American Journal of Epidemiology*, 171: 465-478.
- Jordà, Ò., Singh, S. R., and Taylor, A. M. (2020). *Longer-run Economic Consequences of Pandemics*. Working Paper, National Bureau of Economic Research.
- Lemaitre, M., & Carrat, F. (2010). Comparative age distribution of influenza morbidity and mortality during seasonal influenza epidemics and the 2009 H1N1 pandemic. *BMC infectious diseases*, 10: 1-5.
- Lemaitre, M., Carrat, F., Rey, G., Miller, M., Simonsen, L., and Viboud, C. (2012, 9). Mortality Burden of the 2009 A/H1N1 Influenza Pandemic in France: Comparison to Seasonal Influenza and the A/H3N2 Pandemic. *PLoS ONE*, 7: e45051.

- MacCarthy, F. D. (1969). Hong Kong Flu. *The British Medical Journal*, 1: 182-182.
- McCullers, J. A. (2013). Do specific virus–bacteria pairings drive clinical outcomes of pneumonia? *Clinical Microbiology and Infection*, 19: 113-118.
- Meslé, F. (2010). A spectacular decline in influenza mortality: the role of vaccination. *Population Societies*, 470: 1-4.
- Metersky, M. L., Masterton, R. G., Lode, H., File, T. M., and Babinchak, T. (2012). Epidemiology, microbiology, and treatment considerations for bacterial pneumonia complicating influenza. *International Journal of Infectious Diseases*, 16: e321–e331.
- Morens, D. M., Taubenberger, J. K., and Fauci, A. S. (2009). The Persistent Legacy of the 1918 Influenza Virus. *New England Journal of Medicine*, 361, 225-229.
- Morse, S. S. (2007). Pandemic influenza: Studying the lessons of history. *Proceedings of the National Academy of Sciences*, 104: 7313-7314.
- Piehl, A. M., Cooper, S. J., Braga, A. A. and Kennedy, D. M. (2003). Testing for structural breaks in the evaluation of programs. *Review of Economics and Statistics*, 85(3): 550-558.
- Peretti-Watel, P., Seror, V., Cortaredona, S., Launay, O., Raude, J., Verger, P., Fressard, L., Beck, F., Legleye, S., L’Haridon, O., Léger, D. and Ward, J. K. (2020). A future vaccination campaign against COVID-19 at risk of vaccine hesitancy and politicisation. *The Lancet Infectious Diseases*, 20: 769-770.
- Peretti-Watel, P., Verger, P., Raude, J., Constant, A., Gautier, A., Jestin, C., and Beck, F. (2013). Dramatic change in public attitudes towards vaccination during the 2009 influenza A (H1N1) pandemic in France. *Eurosurveillance*, 18: 20623.
- Quandt, R. E. (1960). Tests of the hypothesis that a linear regression system obeys two separate regimes. *Journal of the American statistical Association*, 55: 324-330.
- Rizzi, S., Sjøgaard, J., & Vaupel, J. W. (2021). High excess deaths in Sweden during the first wave of COVID-19: Policy deficiencies or ‘dry tinder’? *Scandinavian Journal of Public Health*, 50: 33-37.
- Rizzo, C., Bella, A., Viboud, C., Simonsen, L., Miller, M. A., Rota, M. C., Salmaso, S. and degli Atti, M. L. (2007). Trends for Influenza-related Deaths during Pandemic and Epidemic Seasons, Italy, 1969–2001. *Emerging Infectious Diseases*, 13: 694-699.
- Roux, A., Canet, E., Valade, S., Gangneux-Robert, F., Hamane, S., Lafabrie, A., Maubon, D., Debourgogne A., Le Gal, S., Dalle, F., Leterrier, M., Toubas, D., Pomares C., Bellanger, A. P. and Azoulay, É. (2014). iPneumocystis jirovecii/iPneumonia in Patients with or without AIDS, France. *Emerging Infectious Diseases*, 20: 1490-1497.
- Ruhm, C. J. (2000). Are Recessions Good for Your Health? *The Quarterly Journal of Economics*, 115: 617-650.
- Sands, P., Mundaca-Shah, C., and Dzau, V. J. (2016). The Neglected Dimension of Global Security — A Framework for Countering Infectious–Disease Crises. *New England Journal of Medicine*, 374: 1281-1287.

- Schwarzinger, M., Flicoteaux, R., Cortarenoda, S., Obadia, Y., and Moatti, J.-P. (2010). Low Acceptability of A/H1N1 Pandemic Vaccination in French Adult Population: Did Public Health Policy Fuel Public Dissonance? *PLoS ONE*, 5: e10199.
- Shrestha, S., Foxman, B., Berus, J., Van Panhuis, W. G., Steiner, C., Viboud, C., and Rohani, P. (2015). The role of influenza in the epidemiology of pneumonia. *Scientific reports*, 5: 1-13.
- Sohier, R., & Henry, M. (1969). Epidemiological data on Hong Kong influenza in France. *Bulletin of the World Health Organization*, 41: 402-404.
- Toscani, L., Gauthey, L., and Robert, C. (2001). Prévention de la morbidité et mortalité grippale: stratégies d'action. *Forum Médical Suisse – Swiss Medical Forum*.
- Tyrrell, D. A., Buckland, R., Rubenstein, D., and Sharpe, D. M. (1970). Vaccination against Hong Kong Influenza in Britain, 1968-9: A Report to the Medical Research Council Committee on Influenza and Other Respiratory Virus Vaccines. *The Journal of Hygiene*, 68: 359-368.
- Vagneron, F. (2015, 6). Surveiller et s'unir ? *Revue d'anthropologie des connaissances*, 9(2): 139-162.
- Valleron, A.-J., Cori, A., Valtat, S., Meurisse, S., Carrat, F., & Boëlle, P.-Y. (2010, 4). Transmissibility and geographic spread of the 1889 influenza pandemic. *Proceedings of the National Academy of Sciences*, 107: 8778-8781.
- Vaux, S., Van Cauteren, D., Guthmann, J.-P., Le Strat, Y., Vaillant, V., Valk, H., & Lévy-Bruhl, D. (2011). Influenza vaccination coverage against seasonal and pandemic influenza and their determinants in France: a cross-sectional survey. *BMC public health*, 11: 1-9.
- Viboud, C., Grais, R. F., Lafont, B. A., Miller, M. A., & Simonsen, L. (2005, 7). Multinational Impact of the 1968 Hong Kong Influenza Pandemic: Evidence for a Smoldering Pandemic. *The Journal of Infectious Diseases*, 192: 233-248.
- Ward, J. K., Peretti-Watel, P., Bocquier, A., Seror, V., & Verger, P. (2019). Vaccine hesitancy and coercion: all eyes on France. *Nature immunology*, 20: 1257-1259.


Appendix

Figure A1. Death rates of flu by departments Ain to Gard (deaths per 100,000 population).




Notes: This figure plots flu mortality rates by year and department (*Ain* to *Gard*) for the period 1950-2015. The red lines provide estimates of mortality trends for the periods 1950-1970 and 1970-2015 using Poisson model. Hong Kong flu (1968-1970) is shown in grey.

Figure A2. Death rates of flu by departments Haute-Garonne to Oise (deaths per 100,000 population).




Notes: This figure plots flu mortality rates by year and department (Haute-Garonne to Oise) for the period 1950-2015. The red lines provide estimates of mortality trends for the periods 1950-1970 and 1970-2015 using Poisson model. Hong Kong flu (1968-1970) is shown in grey.

Figure A3. Death rates of flu by departments Orne to *Territoire de Belfort* (deaths per 100,000 population).




Notes: This figure plots flu mortality rates by year and department (Haute-Garonne to Oise) for the period 1950-2015. The red lines provide estimates of mortality trends for the periods 1950-1970 and 1970-2015 using Poisson model. Hong Kong flu (1968-1970) is shown in grey.

Figure A4. Death rates of flu by region, with and without 1957 and 1968-1969 pandemics (deaths per 100,000 population).




Notes: This figure plots flu mortality rates by year and region for the period 1950-2015. The red lines provide estimates of mortality trends for the periods 1950-1970 and 1970-2015 using Poisson model. Hong Kong flu (1968-1970) is shown in grey. The blue lines provide estimates of mortality trends for the periods 1950-1970 and 1970-2015 using Poisson model without 1957 and 1968-1969 pandemics.

Figure A5. Death rates of flu by age groups, with and without 1957 and 1968-1969 pandemics.



Notes: This figure plots death rates of flu by year and age group for the period 1950-2015. The red lines provide estimates of mortality trends for the periods 1950-1970 and 1970-2015 using Poisson model. Hong Kong flu (1968-1970) is shown in grey. The blue lines provide estimates of mortality trends for the periods 1950-1970 and 1970-2015 using Poisson model without 1957 and 1968-1969 pandemics.

Figure A6: Flu death rate and vaccination in France from 1950 to 2015 (deaths per 100,000 population).



Notes: This figure plots flu death rate (in black) as well as vaccination rate (in red and green) by year for the period 1950-2015. Vaccination rate before 1977 is estimated according to 4 scenarios: a linear trend (green line), and 3 exponential trends (in red) with vaccination rate equals to 10%, 5% and 2% in 1970. Hong Kong flu (1968-1970) is shown in grey. Vaccination rate from 1977 onwards comes from http://www.infectiologie.com/UserFiles/File/medias/enseignement/du-lyon/2014-DUCIV-Lyon-Escuret_vaccin_grippe.pdf (page 79) and <http://www.grippe-geig.com/couverture-vaccinale.html>.

Figure A7: Evidence from newspapers (12/05/1969; 12/14/1969; 01/07/1970) – France Soir, Paris Presse, L'intransigeant.

LE DOSSIER COMPLET DE LA GRIPPE

« Envoyez vite une standardiste, j'ai du prendre sa place » demande le P.-D.G.

Robert Boulvin (ministre de la Santé publique et de la Sécurité sociale)

« Je suis un exemple vivant de l'utilité du vaccin »

Portrait du virus : une sphère minuscule hérissée de piquants

Europe : 1.000 morts en Italie 800 cercueils en instance d'incinération à Berlin

25 % d'absentéisme

Beux sortes de vaccins

Deux autres médicaments

Opinionnaire aux P.T.T.

Le virus A 2 de Hong-Kong a provoqué sa première tragédie : à Annappes, près de Lille, un bébé de deux mois, le fils de Mme Mande, est mort hier, victime de la grippe.

L'épidémie qui vient d'atteindre le Nord a mis un mois pour franchir les Pyrénées

« La deuxième vague de cette épidémie venue de l'Inde et du Japon en 1968 en passant par les Etats-Unis où elle a fait 20.000 morts l'hiver dernier, surprend on l'attendait en janvier 1970. Elle a fiévre : plus que la maladie qui n'est pas considérée encore comme vraiment grave à l'Institut Pasteur, à Paris, c'est la psychose de l'épidémie qui crée des difficultés.

« Hier, on s'est précipité dans les pharmacies pour acheter des vaccins anti-grippe. Les stocks ont été dévalisés : « J'ai reçu trente coups de téléphone avant l'ouverture du magasin », dit un pharmacien de la place d'Italie. « J'ai vendu vingt-cinq vaccins par heure », dit un autre de la place Wagram. « J'en avais plein mon réfrigérateur le matin. Je n'en ai plus ce soir », ajoute un troisième pharmacien de la care Montparnasse.

A l'Institut Pasteur, on insiste particulièrement sur les signes de la grippe de Hong-Kong à ne pas confondre avec la grippe habituelle de nos hivers (qui fait tout de même 12.000 morts par an) : maux de tête violents, abattement brutal, courbatures, claquements de dents, forte fièvre de 39°, 40°, qui laisse une grande fatigue. La grippe de Hong-Kong est très contagieuse.

Laboratoires impuissants

LA PROTECTION : elle est efficace dans 80 % des cas, avec un des trois vaccins anti-grippe qui répondent aux normes du laboratoire national du contrôle de la santé (vaccin Pasteur, vaccin Merieux, vaccin Dufar). Ils sont conseillés aux sujets « fragiles » : personnes âgées de plus de 65 ans et malades atteints d'affections chroniques) et dans les services collectifs (administrations, crèches).

LE VACCIN : c'est une ampoule seringue de 10 F, à injection sous-cutanée, pratiquée par le médecin traitant ou dans les dispensaires sous l'ampoule. Pour les adultes : une injection suffit avec un rappel chaque année. Pour les enfants de 5 à 10 ans : deux injections à trois semaines d'intervalle. La vaccination n'est pas conseillée pour les enfants plus jeunes sauf « ils sont fragiles. L'immunité commence de quinze jours à trois semaines après la piqûre.

LES PROVISIONS : elles ne sont pas importantes. Les pharmacies assaillies dans l'après-midi ont téléphoné le soir à leurs grossistes qui ont appelé les laboratoires... qui sont impuissants.

« La vaccination aurait dû s'écarter, dit-on à l'Institut Merieux. Les besoins ont dépassé la possibilité de fabrication des vaccins. Nous n'en avons presque plus. »

L'Institut Pasteur a une réserve de 200.000 vaccins. Il en sort à peu près 100.000 par mois, sans pouvoir pousser vraiment la cadence, « car, dit un médecin, le vaccin... »

Très nuageux demain vendredi, prévoit la Météo nationale


La grippe tue les Français le plus

GRIPPE : Services de garde des pharmacies renforcés dimanche

Ces mortels chez des grippe souffrent de troubles vocodiers


Médecins sur le ligne 87

Figure A8. Death rates of flu (in log) for the older age groups in France, 1980 – 2015.




Notes: This figure reproduces Figure A3 and zooms on the period 1980-2015 for 65 and more age groups.

Figure A9a. Pneumonia death rates by region (deaths per 100,000 population).




Notes: This figure plots pneumonia mortality rates by year and region for the period 1950-2015. The red lines provide estimates of mortality trends for the periods 1950-1970 and 1970-2015 using Poisson model. Hong Kong flu (1968-1970) is shown in grey.

Figure A9b. Pneumonia death rates by age (deaths per 100,000 population).




Notes: This figure plots pneumonia mortality rates by year and age group for the period 1950-2015. The red lines provide estimates of mortality trends for the periods 1950-1970 and 1970-2015 using Poisson model. Hong Kong flu (1968-1970) is shown in grey.

Figure A10. All cause death rates by region (deaths per 100,000 population).



Notes: This figure plots all cause mortality rates by year and region for the period 1950-2015. The red lines provide estimates of mortality trends for the periods 1950-1970 and 1970-2015 using Poisson model. Hong Kong flu (1968-1970) is shown in grey.

Figure A11. Excess deaths in France, 1950-2015.



Notes: This figure plots excess deaths caused by flu. We estimate our model presented in equation (1) over the entire period rather than year by year and define excess deaths as the difference between estimated and observed deaths caused by flu.